





Annual Report of the MALAYSIAN PSORIASIS REGISTRY 2007-2011

Editors:

Azura Mohd Affandi Fatimah `Afifah Alias Asmah Johar Roshidah Baba

With contribution from: Tassha Hilda Adnan

Ministry of Health Malaysia

NATIONAL DERMATOLOGY REGISTRY (DermReg)

Annual Report of the MALAYSIAN PSORIASIS REGISTRY

2007 - 2011

Editors:

Azura Mohd Affandi Fatimah `Afifah Alias Asmah Johar Roshidah Baba

With contribution from:

Tassha Hilda Adnan

Jointly published by:

National Dermatology Registry (DermReg), Malaysia, and Clinical Research Centre (CRC), Ministry of Health, Malaysia

Contact:

National Dermatology Registry Department of Dermatology Hospital Kuala Lumpur Jalan Pahang 50586 Kuala Lumpur

Tel: 03-2615 5225 / 03-2615 5555 Ext 5225

Fax: 03-2698 5927

Email: dermreg@acrm.org.my

Website: www.acrm.org.my/dermreg/

Disclaimer

The data reported here have been supplied by DermReg. The interpretation and reporting of these data are the responsibility of the Editor and in no way should be seen as an official policy or interpretation of the DermReg. This report is copyrighted. However it can be freely reproduced without the permission of the DermReg. Acknowledgement would be appreciated.

Suggested citation

The suggested citation for this report is as follows:

Mohd Affandi A, Alias FA, Johar A, Baba R. Annual Report of the Malaysian Psoriasis Registry 2007-2011, Kuala Lumpur, Malaysia 2013

Electronic version

Electronic version of this report can be downloaded at http://www.acrm.org.my/dermreg





ACKNOWLEDGMENT

We would like to thank the Director General of Health, Malaysia for permission to publish this report.

The National Dermatology Registry would like to give its appreciation to everyone who has helped in making this report possible.

We would also like to thank the following for their contribution and support:

- All the doctors, allied health personnel and clerical staff in the participating centres
- The Ministry of Health, Malaysia
- The Dermatological Society of Malaysia
- Clinical Research Centre (CRC), Ministry of Health, Malaysia
- Altus Solutions Sdn Bhd

CONTENTS

| ACKNOWLEDGEMENTS | i |
|---|--|
| CONTENTS | ii |
| LIST OF TABLES | iii |
| LIST OF FIGURES | iv |
| ABBREVIATIONS | v |
| ABOUT DermReg Introduction Objectives Organization Sponsors Governance Board Steering Committee Registry Coordinating Centre Source Data Providers Official website | vi vi vii viii ix x xi |
| ABOUT MPR Introduction Objectives Scope of MPR | xiii xiii xiv |
| EXECUTIVE SUMMARY | 1 |
| CHAPTER 1: STOCK AND FLOW | 3 |
| CHAPTER 2: CHARACTERISTICS OF PATIENTS | 7 |
| CHAPTER 3: MEDICAL HISTORY | 9 |
| CHAPTER 4: COMORBIDITIES | 13 |
| CHAPTER 5: CLINICAL PRESENTATION | 17 |
| CHAPTER 6: TREATMENT | 23 |
| CHAPTER 7: QUALITY OF LIFE | 27 |
| CHAPTER 8: OUTCOMES | 33 |
| APPENDIX A: CASE REPORT FORM Case Report Form Biologic Treatment Initiation Form Biologic Treatment Follow Up Form | 39 43 45 |
| APPENDIX B: DATA MANAGEMENT | 47 |
| APPENDIX C: STATISTICAL METHODS | 53 |
| APPENDIX D: PARTICIPATING CENTRE DIRECTORY | 57 |

LIST OF TABLES

| Table 1.1 Number of psoriasis patients notified from each participating centre | 6 |
|--|----|
| Table 1.2 Distribution of psoriasis patients according to the number of notifications | 6 |
| Table 2.1 Patient demographics | 8 |
| Table 3.1 Positive family history of psoriasis and its relationship with age of onset | 11 |
| Table 3.2 Family members with psoriasis | 11 |
| Table 3.3 Aggravating factors of psoriasis | 12 |
| Table 3.4 Infections which aggravated psoriasis | 12 |
| Table 3.5 Drugs which aggravated psoriasis | 13 |
| Table 4.1 Prevalence of comorbidities in adult psoriasis patients aged 18 and above | 16 |
| Table 4.2 Prevalence of comorbidities in psoriasis patients aged below 18 years | 17 |
| Table 4.3 Co-morbidities associated with psoriatic arthritis patients | 17 |
| Table 5.1 Distribution of psoriasis patients according to the type of psoriasis | 21 |
| Table 5.2 Distribution of percentage of body surface area affected in psoriasis patients | 21 |
| Table 5.3 Distribution of severity of body part affected of psoriasis patients | 21 |
| Table 5.4 Distribution of nail involvement in psoriasis patients | 22 |
| Table 5.5 Distribution of nail features in patients with nail involvement | 22 |
| Table 5.6 Distribution of joint disease in psoriasis patients | 22 |
| Table 5.7 Rheumatoid factor results in psoriasis patients with joint disease | 22 |
| Table 5.8 Distribution of types of joint disease | 23 |
| Table 5.9 Symptoms of psoriatic arthritis | 23 |
| Table 5.10 Distribution of types of joint deformities in patients with joint disease | 23 |
| Table 5.11 Factors associated with psoriatic arthritis | 24 |
| Table 6.1 Use of topical therapy in psoriasis patients | 27 |
| Table 6.2 Distribution of types of topical therapy | 27 |
| Table 6.3 Use of phototherapy in psoriasis patients | 27 |
| Table 6.4 Distribution of types of phototherapy | 27 |
| Table 6.5 Use of systemic therapy in psoriasis patients | 28 |
| Table 6.6 Distribution of types of systemic therapy in psoriasis patients | 28 |
| Table 7.1 Responses for DLQI in adult psoriasis patients (age 17 and above) | 31 |
| Table 7.2 Responses for CDLQI in child/adolescent psoriasis patients (aged 5 to 16) | 33 |
| Figure 7.3 Quality of life in children/adolescents with psoriasis | 34 |
| Table 8.1 Distribution of psoriasis patients according to the duration of follow-up | 37 |

LIST OF FIGURES

| Figure 1.1 | Psoriasis patients notified to the MPR | 5 |
|------------|---|----|
| Figure 1.2 | Psoriasis patients notified to the MPR per year | 5 |
| Figure 3.1 | Distribution of age of onset | 11 |
| Figure 7.1 | Quality of life in adult patients with psoriasis | 32 |
| Figure 7.2 | QoL impairment in adults psoriasis patients based on category of DLQI | 32 |
| Figure 7.3 | Quality of life in children/adolescents with psoriasis | 34 |
| Figure 7.4 | QoL impairment in child/adolescent psoriasis patients based on category | 34 |
| | of DLQI | |
| Figure 8.1 | Improvement in the extent of skin lesions | 38 |
| Figure 8.2 | Improvement in total clinical skin scores | 38 |
| Figure 8.3 | Improvement in joint pain | 39 |
| Figure 8.4 | Improvement in DLQI and CDLQI | 39 |

ABBREVIATIONS

BB-UVB Broad-band ultraviolet B

Body mass index BMI Body surface area **BSA**

Child Dermatology Life Quality Index **CDLOI**

CRC Clinical Research Centre

CRF Case report form

DermReg National Dermatology Registry **DLQI** Dermatology Life Quality Index eCRF Electronic case report form DermReg web application eDermReg HLA Human leukocyte antigen

IOR Interquartile range Ministry of Health MOH

Malaysian Psoriasis Registry **MPR**

Not available NA

Narrow-band ultraviolet B **NBUVB**

National Health and Morbidity Survey **NHMS**

PΙ Principal Investigator Psoralen and ultraviolet A **PUVA**

QoL Quality of life

RCC Registry Coordinating Centre

SC Site Coordinator SD Standard deviation **SDP** Sources data providers

ABOUT DermReg

Introduction

DermReg is an ongoing systematic collection, analysis and interpretation of data pertaining to dermatological diseases and services in Malaysia. It is a nationwide project which aims to integrate all dermatological patient registries and databases developed in Malaysia. These registries are essential in the planning, implementation and evaluation of clinical and health services as well as research in dermatology

Objectives of DermReg

General Objective


To establish a nationwide systematic prospective collection of data pertaining to skin diseases and dermatological services, in order to study the natural history, outcome and quality of life issues of skin diseases, as well as the effectiveness, safety and accessibility of various treatment modalities.

Specific Objectives:

- 1. Determine the socio-demographic profile of patients with skin diseases
- 2. Determine the burden of skin diseases in the population
- 3. Describe the natural history of skin diseases
- 4. Identify the potential causal and risk factors of skin diseases
- 5. Describe the clinical manifestation of skin diseases
- 6. Describe the effect of skin diseases on the quality of life
- 7. Determine the efficacy and cost effectiveness of treatment of skin diseases
- 8. Monitor the safety and adverse effects of products and services used in the treatment
- 9. Evaluate accessibility and quality of health services related to skin diseases
- 10. Stimulate and facilitate basic, clinical and epidemiological research on skin diseases

ORGANISATION OF DermReg

The organizational structure of DermReg consists of sponsors, Governance Board, Steering Committee, Sub-committees or Expert Panels, Registry Coordinating Centre, Source Data Providers (SDP) and users.



SPONSORS

The DermReg is sponsored by:

- Ministry of Health, Malaysia 1.
- 2. Clinical Research Centre, Hospital Kuala Lumpur
- 3. The Dermatological Society of Malaysia
- Pharma companies Abbott Malaysia, Leo Pharma Malaysia and Janssen 4. Malaysia

GOVERNANCE BOARD

Governance Board of DermReg is a committee established by the sponsors. Its roles are:

- to ensure that the DermReg stay focused on its objectives
- to ensure its continuing relevance and justification
- 1. Datuk Dr. Roshidah Baba (Chairperson) Head of Dermatological Services and Senior Consultant Dermatologist Department of Dermatology Hospital Melaka
- 2. Dr. Najeeb Ahmad Mohd Safdar President of the Dermatological Society of Malaysia, and Consultant Dermatologist Hospital Tuanku Jaafar, Seremban Negeri Sembilan
- 3. Dr. Steven Chow Kim Weng President of the College of Physicians, Academy of Medicine Malaysia, and Senior Consultant Dermatologist The Skin Centre, Kuala Lumpur
- 4. Dr. Goh Pik Pin Director of the Clinical Research Centre Network Ministry of Health

STEERING COMMITTEE

Steering Committee for Malaysian Psoriasis Registry (MPR)

| No. | Name | Institution |
|-----|--|---------------------------------------|
| 1. | Dr Chang Choong Chor (2007-Jul 2012) Dr. Azura Mohd Affandi (July 2012 – current) | Hospital Kuala Lumpur |
| 2. | Dr. Choon Siew Eng | Hospital Sultanah Aminah, Johor Bahru |
| 3. | Dr. Pubalan Muniandy | Hospital Umum Sarawak |
| 4. | Dr. Tang Jyh Jong | Hospital Permaisuri Bainun, Ipoh |
| 5. | Dr. Chan Lee Chin | Hospital Pulau Pinang |
| 6. | Dr. Najeeb Ahmad Mohd Safdar | Hospital Tuanku Jaafar, Seremban |
| 7. | Dr. Steven Chow Kim Weng | The Skin Clinic, Kuala Lumpur |
| 8. | Dr. Mohd Noh Idris | Klinik Kulit Md Noh, Kuala Lumpur |

REGISTRY COORDINATING CENTRE

The DermReg Registry Coordinating Centre (RCC) is based at the Department of Dermatology, Hospital Kuala Lumpur. It coordinates the data collection among the source data providers, and collaborates with the Clinical Research Centre (CRC) that provides epidemiological and statistical support.

Registry Manager Fatimah 'Afifah Alias

Technical Support Personnel

Epidemiology Officer Dr. Jamaiyah Haniff

Clinical Epidemiology Unit,

CRC

Biostatisticians Ms Tassha Hilda bt Adnan

CRC

Database Administrator Ms Lim Jie Ying

Altus Solutions Sdn Bhd

SOURCE DATA PROVIDERS (SDP)

Source data providers (SDP) are centres that contribute data to the registries.

Source Data Providers for Malaysian Psoriasis Registry (MPR)

| No. | Source Data Provider | Investigator |
|-----|---|--------------------------------|
| 1. | Hospital Kuala Lumpur | Dr. Azura Mohd Affandi |
| 2. | Hospital Pulau Pinang | Dr. Chan Lee Chin |
| 3. | Hospital Sultanah Bahiyah, Alor Setar | Dr. Mani Mala a/p T. Manikam |
| 4. | Hospital Tuanku Fauziah, Perlis | Dr. Sharifah Farihah Syed Abas |
| 5. | Hospital Sultanah Fatimah, Muar | Dr. Siti Khadijah Abdul Wahid |
| 6. | Hospital Tuanku Jaafar, Seremban | Dr. Najeeb Ahmad Mohd Safdar |
| 7. | Hospital Queen Elizabeth, Kota Kinabalu | Dr. Zaigham Mahmood |
| 8. | Hospital Sungai Buloh | Dr. Azahzuddin Hamzah |
| 9. | Hospital Tengku Ampuan Afzan, Kuantan | Dr. Abu Razak Yusof |
| 10. | Hospital Permaisuri Bainun, Ipoh | Dr. Tang Jyh Jong |
| 11. | Hospital Umum Sarawak, Kuching | Dr. Pubalan Muniandy |
| 12. | Hospital Tengku Ampuan Rahimah, Klang | Dr. Ng Ting Guan |
| 13. | Hospital Melaka | Dr. Che Salmi Yusoff |
| 14. | Prince Court Medical Centre | Dr.Gangaram Hemandas |
| 15. | Gleneagles Intan Medical Centre | Dr. Chang Choong Chor |
| 16. | Hospital Sultanah Aminah, Johor Bahru | Dr. Choon Siew Eng |
| 17. | Hospital Universiti Kebangsaan Malaysia | Dr. Mazlin Mohd Baseri |
| 18. | Pusat Perubatan Universiti Malaya | Dr. Wong Su Ming |

OFFICIAL WEBSITE OF DermReg

http://www.acrm.org.my/dermreg/



Visitor number: 007548

Copyright © 2009 by Altus Solutions 5dn Bhd - Best Viewed in 1024 x 768 resolution | Last updated: 10/05/2012

ABOUT MALAYSIAN PSORIASIS REGISTRY (MPR)

Introduction

Psoriasis is a common skin disease, characterized by inflamed scaly patches and plaques. It runs a chronic relapsing course with variable degree of severity, and causes significant physical, psychosocial and economic impact on the patient. Being incurable, it may lead to poor patient compliance especially in treatment which will further compromise the overall management of the disease.

The Malaysian Psoriasis Registry (MPR) is a skin disease clinical registry. It is a prospective, ongoing systematic collection of data pertaining to patients who have psoriasis. The main reason for setting up a psoriasis registry is to have more accurate data on the various aspects of psoriasis in Malaysia. This would help in assessing the true magnitude of the problem in Malaysia, including the demographic data, types of psoriasis, its severity, aggravating factors, any associated joint and nail involvement and the various types of therapies commonly used. Having a psoriasis registry would also help in research work and more importantly in improving the overall management of the patients.

Preliminary work on the MPR started in 1998 by a group of dermatologists, which culminated in the First Malaysian Psoriasis Symposium on the 17th May 1998. This registry consists of information on patients with psoriasis in Malaysia and is under the umbrella of the National Dermatology Registry (DermReg). A case report form was developed and data collection started as a pilot project in March 2000. A preliminary report of the registry (March 2000 to July 2005) was published in the Malaysian Journal of Dermatology in the August 2005 issue.

In 2007, MPR was extensively revised under the guidance of CRC and with the financial support from MOH. A new case report form was introduced and a new centralised electronic database with web application was established to facilitate multi-centre data collection. Preliminary report of the newly revised MPR was published in the Medical Journal of Malaysia in September 2008. The First Annual Report of MPR 2007-2008 was published in the following year.

Objectives

The MPR has the following objectives:

> Primary objective:

To obtain more accurate data on various aspects of psoriasis in Malaysia.

- > Secondary objectives:
 - To determine the socio-demographic profiles of patients with psoriasis.
 - To determine the disease burden attributed to psoriasis.
 - To provide information for planning of medical services, facilities, manpower and training related to the management of psoriasis.
 - 4. To stimulate and facilitate research on psoriasis and its management.

Scope of MPR

The MPR is intended to be a truly national population based disease and treatment registry. Hence it seeks the participation of all providers of dermatological services in both the public and private sectors in Malaysia.

The MPR collects:

- Demographic data
- ➤ Clinical data including patients' history and clinical examination findings
- Quality of life measure i.e. Dermatology Life Quality Index (DLQI)
- > Modalities of treatment used

Outcomes of interest include:

- > Course of the disease
- ➤ How the disease affects quality of life
- > Disease improvement with treatment
- ➤ Association with any other diseases

Inclusion criteria:

1. All patients who are clinically diagnosed to have psoriasis by a registered dermatologist or by a medical practitioner under the supervision of a dermatologist are included. Confirmation of diagnosis by histopathologic examination is optional.

Exclusion criteria:

Patients whose diagnosis is in doubt are excluded.

EXECUTIVE SUMMARY

Stock and Flow

During the period from October 2007 to December 2011, a total of 6,985 patients with psoriasis from 19 dermatology centres (15 governments, 2 private centres and 2 university hospitals) were notified to the registry.

Demographic Characteristics of Patients

Male-to-female ratio was 1.3:1. Ethnic distribution: Malay 50.2%, Chinese 22.3%, Indian 18.1%, other ethnic groups 9.3% and Orang Asli 0.1%. Mean age at notification was 43 \pm 17.33 years (range 0 - 97 years). Most patients (99.1%) were Malaysian citizens.

Medical History

Mean age of onset of psoriasis was 33.3 ± 16.7 years (range 0 - 85 years). Family history of psoriasis was present in 20.1% of the patients. Positive family history was more common among patients with younger onset (aged 40 and below) compared to those with later onset of disease: 15.5% vs 4.5%. Among those who had positive family history, family members affected were either of their parents in 41.8%, siblings in 36.7% and children in 10.6%.

51.1% of the patients reported one or multiple factors which aggravated their psoriasis. The commonest aggravating factors were stress (67.0%), sunlight (35.9%) and infection (17.4%).

Comorbidities

In adult psoriasis patients aged 18 and above, the common comorbidities were overweight 33.6%, obesity 22.5%, hypertension 25.4%, diabetes mellitus 17.8%, hyperlipidaemia 16.6%, and ischaemic heart disease 5.6%. In patients aged below 18, the commonest comorbidity was overweight or obesity (27.9 %), bronchial asthma (2.2%) followed by diabetes mellitus (0.4%) down syndrome (0.4%) and congenital heart disease (0.4%).

Clinical Presentation

The commonest clinical type of psoriasis was plaque psoriasis (88.9%). This was followed by guttate psoriasis (4.5%), erythrodermic psoriasis (2.1%), pustular psoriasis (1.4%) and flexural psoriasis (0.5%). The majority of patients (75.9%) had body surface area involvement of 10% or less.

Psoriatic arthropathy was reported in 15.0% of patients. The commonest psoriatic arthropathy oligo/monoarthropathy (49.1%) followed by rheumatoid-like symmetrical polyarthropathy (36.4%) and distal hand joints arthropathy (31.4%). Spondylitis/sacroilitis accounted for only 11.1% of cases and arthritis mutilans in 3.4% of patients.

About two-third (59.8%) of patients had nail changes associated with psoriasis. Among patients who had nail disease, pitting was commonest (72.5%), followed by onycholysis (50.3%), discoloration (36.1%) and subungual hyperkeratosis (15.0%). Total nail dystrophy was found in 5.4% of patients with nail disease.

Treatments received in the past 6 months

72.6% of the patients were on topical treatment only. Topical steroid was the commonest prescribed (85.4%), followed by tar preparations in 80.7% and emollients in 74.7% of patients. 4.1% of patients received phototherapy. Of the patients who had phototherapy, narrowband UVB (NBUVB) was the commonest used (72.8%). Oral psoralen and ultraviolet A (PUVA) was used in 4.0% of patients only. Systemic therapy was given in 19.6% of patients. The most frequently used systemic therapy was methotrexate (67.8%), followed by acitretin (20.8%), systemic corticosteroids (6.3%), sulphasalazine (6.2%), cyclosporine (4.5%), biologics (1.9%) and hydroxyurea (1.2%).

Quality of Life

Measurement of quality of life using Dermatology Life Quality Index (DLQI) or Child DLQI (CDLQI) was performed in 4,295 adult patients (aged 17 and above) and 245 children/adolescent patients (aged 5 to 16). The mean DLQI score was 8.5 ± 6.4 for adult patients and the mean CDLQI was 7.6 ± 5.5 for children/adolescent patients.

CHAPTER 1

STOCK AND FLOW

During the period from October 2007 to December 2011, a total of 6,985 patients were notified to the registry. The number of notified patients gradually increased throughout the period (Figure 1.1). The number of patients notified per year is illustrated in Figure 1.2.

A total of 19 dermatology centres (15 government, 2 private centres and 2 university hospital) participated in the MPR. Department of Dermatology, Hospital Kuala Lumpur notified the highest number of patients. This was followed by Hospital Pulau Pinang and Hospital Tengku Ampuan Rahimah, Klang (Table 1.1).

The majority of the patients (72.2%) were notified once. A second notification during subsequent follow-up visit was also received in 1943 (27.8%) patients. Out of these patients, 1166 (16.7%) had one follow-up notification, 448 (6.4%) had two follow-up notifications, and 329 (4.6%) had more than two follow-up notifications (**Table 1.2**).




Figure 1.1 Psoriasis patients notified to the MPR




Figure 1.2 Psoriasis patients notified to the MPR per year

Table 1.1 Number of psoriasis patients notified from each participating centre

| | No. of patients notified | | | | | |
|---|--------------------------|------|------|------|------|-------|
| | 2007 | 2008 | 2009 | 2010 | 2011 | Total |
| Hospital Kuala Lumpur | 83 | 264 | 338 | 236 | 125 | 1046 |
| Hospital Pulau Pinang | 21 | 92 | 293 | 158 | 232 | 796 |
| Hospital Tengku Ampuan Rahimah, Klang | 0 | 83 | 199 | 303 | 148 | 733 |
| Hospital Melaka | 0 | 0 | 97 | 302 | 257 | 656 |
| Hospital Umum Sarawak, Kuching | 6 | 199 | 150 | 113 | 162 | 630 |
| Hospital Sultanah Bahiyah, Alor Setar | 109 | 224 | 96 | 78 | 63 | 570 |
| Hospital Queen Elizabeth, Kota Kinabalu | 20 | 107 | 131 | 145 | 141 | 544 |
| Hospital Raja Permaisuri Bainun, Ipoh | 64 | 60 | 151 | 62 | 141 | 478 |
| Hospital Tengku Ampuan Afzan, Kuantan | 0 | 50 | 56 | 144 | 146 | 396 |
| Hospital Sultanah Fatimah, Muar | 4 | 44 | 29 | 55 | 189 | 321 |
| Hospital Sultanah Aminah, Johor Bahru | 0 | 38 | 148 | 69 | 66 | 321 |
| Hospital Tuanku Fauziah, Kangar | 1 | 33 | 56 | 69 | 55 | 214 |
| Hospital Tuanku Jaafar, Seremban | 0 | 44 | 0 | 38 | 50 | 132 |
| Hospital Sungai Buloh | 8 | 29 | 2 | 0 | 0 | 39 |
| Prince Court Medical Centre | 0 | 0 | 6 | 18 | 3 | 27 |
| Gleneagles Medical Centre | 0 | 16 | 6 | 0 | 0 | 22 |
| UKMMC | 0 | 0 | 0 | 17 | 0 | 17 |
| Total | 316 | 1283 | 1758 | 1807 | 1821 | 6985 |

 Table 1.2
 Distribution of psoriasis patients according to the number of notifications

| Year | No. | % |
|---------------------------------------|--------|------|
| Entry notification | 5, 042 | 72.2 |
| Entry and one follow-up notifications | 1,166 | 16.7 |
| Entry and 2 follow-up notifications | 448 | 6.4 |
| Entry and 3 follow-up notifications | 175 | 2.5 |
| Entry and 4 follow-up notifications | 99 | 1.4 |
| Entry and 5 follow-up notifications | 38 | 0.5 |
| Entry and 6 follow-up notifications | 16 | 0.2 |
| Entry and 7 follow-up notifications | 1 | 0.0 |
| Total | 6985 | 100 |

CHAPTER 2

CHARACTERISTICS OF PATIENTS

There were more males than females (56.5% and 43.5% respectively), with a male to female ratio of 1.3:1. Malays comprised the majority of patients (50.2%), followed by Chinese (22.3) Indians (18.1%), other ethnic groups (9.3%) and Orang Asli (0.1%).

The mean age at presentation to the clinic was 43.0 ± 17.33 years with a range from 0 to 97 years. The majority were married (67.0%), 28.5% were single, and the rest, either divorced or widowed (Table 2.1).

Table 2.1 Patient demographics

| Patient characteristics | | No | % |
|-----------------------------|-------------------|----------------------------|------|
| Gender | Male | 3, 948 | 56.5 |
| | Female | 3, 037 | 43.5 |
| | Malay | 3, 506 | 50.2 |
| | Chinese | 1, 557 | 22.3 |
| | Indian | 1, 262 | 18.1 |
| | Orang Asli | 9 | 0.1 |
| | Others | 651 | 9.3 |
| | Malaysian | 6, 919 | 99.1 |
| | Non Malaysian | 58 | 1.0 |
| | Single | 1, 991 | 28.5 |
| | Married | 4, 678 | 67.0 |
| | Divorced | 56 | 0.8 |
| | Widowed | 139 | 2.0 |
| | NA | 121 | 1.8 |
| Age at notification (years) | Mean ± SD (Range) | $43.0 \pm 17.3 \ (0 - 97)$ | |

CHAPTER 3

MEDICAL HISTORY

Onset of Psoriasis

Psoriasis may first appear at any age. In the MPR, 67.0% of patients had first symptoms of psoriasis by the age of 40. The mean age of onset in our cohort was 33.3 ± 16.7 years with a wide range from 0 to 85 years (Figure 3.1). The mean interval between onset (as reported by patient) and diagnosis (first diagnosed by physician) was 2.12 ± 4.5 years.

Family History

Psoriasis is a skin disorder with a polygenic mode of inheritance. In our registry, about onefifth (20.1%) of patients had at least one family member who has psoriasis (**Table 3.1**). Of those with a positive family history, 41.8% had either of their parents affected. Siblings were affected in 36.7% and children in 10.6% (Table 3.2). More patients with positive family history were observed among those with a younger onset of disease (aged 40 and below) compared to those with later onset of disease: 15.5% vs 4.5% (**Table 3.1**).

Aggravating factors of psoriasis

51.1% of patients reported one or multiple factors which worsened their psoriasis. Stress was the commonest aggravating factor (67.0%), followed by sunlight (35.9%) and infection (17.4%). Other identified aggravating factors included trauma (7.7%), smoking (6.1%), drugs (5.5%), alcohol (3.5%), pregnancy (2.7%) and topical treatment (1.4%) (**Table 3.3**).

Analyzing the subgroup of patients who reported infection as an aggravating factor, upper respiratory tract infection (13.2%) appeared to be the commonest infective trigger (**Table** 3.4). Common medications found to aggravate psoriasis were withdrawal of systemic steroids (30.6%), beta blocker (26.4%), traditional medication/ homeopathy (10.7%), non-steroidal anti-inflammatory drugs (9.1%) and antibiotics (8.3%) (**Table 3.5**).




Figure 3.1 Distribution of age of onset

Table 3.1 Positive family history of psoriasis and its relationship with age of onset

| | | Overall - | | Aş | ge of onse | t of psoriasi | is |
|----------------|-----|-----------|------|--------|------------|---------------|-------|
| | | | | 40 & b | elow | Abo | ve 40 |
| | | No. | % | N | % | N | % |
| Family members | Yes | 1, 390 | 20.1 | 1, 067 | 15.5 | 310 | 4.5 |
| with psoriasis | No | 5, 533 | 79.9 | 3, 537 | 51.5 | 1, 958 | 28.5 |
| Total | | 6, 923 | 100 | 4, 604 | 67.0 | 2, 268 | 33.0 |

Table 3.2 Family members with psoriasis

| Family member (one or multiple) | No. | % |
|---------------------------------|-----|------|
| Father | 365 | 26.3 |
| Mother | 215 | 15.5 |
| Sibling(s) | 510 | 36.7 |
| Children | 148 | 10.6 |
| Others | 349 | 25.1 |

Aggravating factors of psoriasis **Table 3.3**

| Aggravating factors (one or multiple) | No. | % |
|---------------------------------------|--------|------|
| Stress | 2, 358 | 67.0 |
| Sunlight | 1, 262 | 35.9 |
| Infection | 613 | 17.4 |
| Trauma | 272 | 7.7 |
| Smoking | 214 | 6.1 |
| Drugs | 194 | 5.5 |
| Alcohol | 122 | 3.5 |
| Pregnancy | 96 | 2.7 |
| Topical treatment | 50 | 1.4 |

Infections which aggravated psoriasis **Table 3.4**

| Infection | No. | % |
|-----------------------------------|-----|------|
| Upper respiratory tract infection | 81 | 13.2 |
| Febrile illness | 23 | 3.8 |
| HIV | 3 | 0.5 |
| Chickenpox | 3 | 0.5 |
| Viral infection | 2 | 0.3 |
| Dengue fever | 2 | 0.3 |
| Chikugunya | 1 | 0.2 |
| Boil | 1 | 0.2 |

Drugs which aggravated psoriasis **Table 3.5**

| Drug | N | % |
|--------------------------------|-----|------|
| Systemic steroids (withdrawal) | 37 | 30.6 |
| Beta-blocker | 32 | 26.4 |
| Traditional/ Homeopathy | 13 | 10.7 |
| NSAIDs / analgesics | 11 | 9.2 |
| Antibiotic | 10 | 8.3 |
| Antimalarial drug | 3 | 2.5 |
| ACE inhibitor | 3 | 2.5 |
| Oral contraceptive pill | 2 | 1.7 |
| Topical tar preparation | 2 | 1.7 |
| Sodium valporate | 1 | 0.8 |
| Daivobet | 1 | 0.8 |
| "Gamat" (Sea cucumber extract) | 1 | 0.8 |
| Oral hypoglycemic | 1 | 0.8 |
| Tomoxifen | 1 | 0.8 |
| Paracetamol | 1 | 0.8 |
| Supplements | 1 | 0.8 |
| Unknown | 1 | 0.8 |
| Total | 121 | 100 |

CHAPTER 4

COMORBIDITIES

Patients with psoriasis were found to have a number of other concomitant diseases. As the spectrum of diseases differs among age groups, adult and children/adolescent patients were analysed separately.

In adult psoriasis patients aged 17 and above, 33.6% were overweight and 22.5% were obese. 25.4% patients had hypertension, 17.8% had diabetes mellitus, 16.6% had dyslipidaemia, 5.6% had ischaemic heart disease and 1.5% had previous history of stroke (**Table 4.1**).

In children and adolescents aged below 17 years with psoriasis, the most prevalent comorbidity was overweight or obesity i.e. BMI at or above 85th centile (27.9 %), followed by bronchial asthma (2.2%) and diabetes mellitus (0.4%), Down syndrome (0.4%) and congenital heart disease (0.4%). Other comorbid conditions were much less common (Table **4.2**).

Compared to patients without arthritis, patients with psoriatic arthritis were found to have increased co-morbidities such as diabetes mellitus, hypertension, hyperlipidaemia and obesity (p < 0.001) (**Table 4.3**).

Prevalence of comorbidities in adult psoriasis patients aged 17 and above Table 4.1

| Co-morbidity | No. | % |
|-------------------------|-------------------------|--|
| Overweight | $2,258^{\#}(2,051^{*})$ | 39.2 [#] (33.6 [*]) |
| Obesity | $2,010^{\#}(1,363^{*})$ | 34.9# (22.4*) |
| Hypertension | 1, 634 | 25.2 |
| Diabetes mellitus | 1, 141 | 17.6 |
| Dyslipidaemia | 1, 078 | 16.6 |
| Ischaemic heart disease | 357 | 5.5 |
| Stroke | 99 | 1.5 |

[#] according to BMI classification for adult Asians as stated in the Clinical Practice Guidelines on Management of Obesity 2004, Ministry of Health, Malaysia

^{*} according to the WHO International Classification of BMI.

Table 4.2 Prevalence of comorbidities in psoriasis patients aged below 17 years

| Comorbidity | N | % |
|---|-----|------|
| Overweight or obesity (BMI\ge 85 th centile) | 127 | 27.9 |
| Bronchial asthma | 10 | 2.2 |
| Diabetes mellitus | 2 | 0.4 |
| Hypertension | 1 | 0.2 |
| Hyperlipidaemia | 1 | 0.2 |
| Down syndrome | 2 | 0.4 |
| Thalassemia | 1 | 0.2 |
| Atrial defect | 1 | 0.2 |
| Congenital heart disease | 2 | 0.4 |
| Obstructive sleep apnoea | 1 | 0.2 |

Table 4.3 Co-morbidities associated with psoriatic arthritis patients

| Co-morbidities | Arthritis Absent (n=5837) | | Arthritis Present (n=1034) | | Simple Logistic Regression | | |
|-------------------------|---------------------------------|--------|----------------------------------|--------|----------------------------|--------------|----------------------|
| | n | (%) | n | (%) | Crude OR | (95% CI) | P-value ^a |
| Diabetes Mellitus | 905 | (15.5) | 218 | (21.1) | 1.47 | (1.24, 1.73) | < 0.001 |
| Hypertension | 1274 | (21.8) | 332 | (32.1) | 1.70 | (1.47, 1.96) | < 0.001 |
| Hyperlipidaemia | 807 | (13.8) | 250 | (24.2) | 2.00 | (1.71, 2.35) | < 0.001 |
| Ischaemic heart disease | 288 | (4.9) | 65 | (6.3) | 1.29 | (0.98, 1.70) | 0.073 |
| Cerebrovascular disease | 85 | (1.5) | 13 | (1.2) | 0.86 | (0.48, 1.55) | 0.610 |
| BMI ≥ 30 (obesity WHO) | 1069 | (18.3) | 250 | (24.2) | 1.42 | (1.21, 1.67) | < 0.001 |

^{*}Result was based on available information. Percentage (%) was calculated based on number of cases over total number for each group (absent or present).

^a Wald statistic.

Chapter 4: Comorbidities

CHAPTER 5

CLINICAL PRESENTATION

Plaque psoriasis was the commonest type of psoriasis (88.9%). This was followed by guttate psoriasis (4.5%) and erythrodermic psoriasis (2.1%). Pustular and flexural/inverse psoriasis were much less common, constituting 1.4% and 0.5% respectively (**Table 5.1**).

Majority of our patients had mild to moderate body surface area involvement. 34.8% of our psoriatic patients had <5% BSA affected, while 41.1% had 5-10% of BSA affected. Severe psoriasis with >10% BSA affected occurred in 21.8% patients, while 2.3% had erythrodermic psoriasis, i.e. >90% BSA involved (**Table 5.2**).

A composite clinical scoring system was used to evaluate the severity of psoriatic lesions in five body regions. A score of 0 to 3 was given for each body region according to the degree of erythema, thickness and scaliness of the skin lesions. The total clinical score may range from 0 to 15. Analysis on severity of lesion noted that most of the moderate to severe lesions (score 2 and 3) located on lower limb (37.0%), trunk (33.2%) and upper limb (29.3%). Half of our psoriatic patients (51.3%) did not have any lesion on the face and neck. If present, lesions on face and neck were generally less severe (score 1 or 2) (**Table 5.3**).

Majority of patients with psoriasis had nail involvement (59.8%) (Table 5.4). Among patients who had psoriatic nail disease, most of them had pitting (72.5%). Other common features were onycholysis (50.3%), discoloration (36.1%) and subungual hyperkeratosis (15.0%). Total nail dystrophy was found in 5.4% of patients with nail involvement (**Table 5.5**).

Joint disease related to psoriasis was reported in 15.0% of the patients (Table 5.6). Rheumatoid factor was detected in 1.8% of patients with arthropathy who were tested (**Table** 5.7). The commonest clinical pattern of psoriatic arthropathy was oligo-/monoarthropathy (49.1%). This was followed by rheumatoid-like symmetrical polyarthropathy (36.4%), distal hand joints arthropathy (31.4%), spondylitis (11.1%) and arthritis mutilans (3.4%). Morning stiffness was observed in 31.5% and enthesopathy in 9.9% (**Table 5.8**).

Most of the patients with psoriatic arthropathy experienced joint pain at time of presentation (86.0%) (**Table 5.9**). Joint swelling was present in 33.6%, while joint deformity occurred in 25.0% (**Table 5.9**). The commonest type of joint deformity was swan neck deformity (27.9%). This was followed by Boutonniere deformity (15.4%), fixed flexion deformity (14.7%), distal hand joint deformity (8.1%), rheumatoid arthritis-like (5.1%) and proximal interphalangeal joint deformity (2.9%). Bamboo spine and arthritis mutilans were observed in only 2.2% of patients (Table 5.10).

By using multiple logistic regressions, 10 factors were found to be associated with psoriatic arthritis. These were age (>18 years), age of onset (\le 40 years), duration of disease (> 5 years), female gender, Indian ethnicity, BMI \geq 30, patients with body surface area \geq 10%, patients with total skin score ≥ 10 , presence of nail involvement and DLQI ≥ 10 (**Table 5.11**).

Table 5.1 Distribution of psoriasis patients according to the type of psoriasis

| Type of psoriasis | No. | % |
|---------------------------|--------|------|
| Plaque | 6, 071 | 88.9 |
| Guttate | 306 | 4.5 |
| Erythrodermic | 146 | 2.1 |
| Pustular | 99 | 1.4 |
| Flexural/Inverse | 31 | 0.5 |
| Palmoplantar non-pustular | 10 | 0.1 |
| Others | 167 | 2.4 |
| Total | 6, 830 | 100 |

Table 5.2 Distribution of percentage of body surface area affected in psoriasis patients

| % Body surface area affected (N=5, 455) | No. | % |
|---|--------|------|
| <5% | 1, 900 | 34.8 |
| 5 - 10% | 2, 240 | 41.1 |
| >10% to 90% | 1, 188 | 21.8 |
| >90% | 124 | 2.3 |
| Total | 5, 452 | 100 |

Table 5.3 Distribution of severity of body part affected in psoriasis patients

| | | | | Clini | cal score | | | |
|-------------|-------|------|-------|-------|-----------|------|-----|-----|
| Body part | 0 | | 1 | | 2 | | 3 | |
| _ | No. | % | No. | % | No. | % | No. | % |
| Scalp | 1,418 | 20.7 | 3,537 | 51.7 | 1,567 | 22.9 | 314 | 4.6 |
| Face & neck | 3,472 | 51.3 | 2,742 | 40.5 | 494 | 7.3 | 64 | 0.9 |
| Trunk | 1,807 | 26.6 | 2,738 | 40.3 | 1,949 | 28.7 | 304 | 4.5 |
| Upper limbs | 1,661 | 24.4 | 3,147 | 46.3 | 1,763 | 25.9 | 229 | 3.4 |
| Lower limbs | 1,362 | 20.0 | 2,926 | 43.0 | 2,162 | 31.8 | 356 | 5.2 |

Table 5.4 Distribution of nail involvement in psoriasis patients.

| Nail involvement | No. | % |
|------------------|--------|------|
| Yes | 4,124 | 59.8 |
| No | 2,773 | 40.2 |
| Total | 6, 897 | 100 |

Table 5.5 Distribution of nail features in patients with nail involvement

| Nail features | No. | % |
|--------------------------|-------|------|
| Pitting | 2,989 | 72.5 |
| Onycholysis | 2,076 | 50.3 |
| Discoloration | 1,487 | 36.1 |
| Subungual hyperkeratosis | 619 | 15.0 |
| Total nail dystrophy | 222 | 5.4 |

Distribution of joint disease in psoriasis patients **Table 5.6**

| Joint disease | No. | % |
|---------------|--------|------|
| Yes | 1,034 | 15.0 |
| No | 5,849 | 85.0 |
| Total | 6, 883 | 100 |

Table 5.7 Rheumatoid factor results in psoriasis patients with joint disease

| Rheumatoid factor | No. | % |
|-------------------|-----|------|
| Positive | 17 | 1.8 |
| Negative | 196 | 21.1 |
| Not available | 717 | 77.1 |
| Total | 930 | 100 |

Table 5.8 Distribution of types of joint disease

| Type of joint disease (one or multiple) | No. | % |
|---|-----|------|
| Oligo-/monoarthropathy | 466 | 49.1 |
| Symmetrical polyarthropathy (rheumatoid-like) | 342 | 36.4 |
| Morning stiffness > 30 mins | 294 | 31.4 |
| Distal hand joints arthropathy | 294 | 31.4 |
| Spondylitis | 102 | 11.1 |
| Enthesopathy | 91 | 9.9 |
| Arthritis mutilans | 31 | 3.4 |

Symptoms of psoriatic arthritis **Table 5.9**

| | Status | No. | % |
|-----------------|--------|-----|------|
| Pain | Yes | 852 | 86.0 |
| | No | 139 | 14.0 |
| Joint swelling | Yes | 332 | 33.6 |
| | No | 655 | 66.4 |
| Joint deformity | Yes | 246 | 25.0 |
| | No | 739 | 75.0 |

Distribution of type of joint deformities in patients with joint disease **Table 5.10**

| Type of joint deformity | No. | % |
|--|-----|------|
| Swan neck deformity | 38 | 27.9 |
| Boutonniere deformity | 21 | 15.4 |
| Fixed-flexion deformity | 20 | 14.7 |
| Distal hand joint deformity | 11 | 8.1 |
| Rheumatoid arthritis-like | 7 | 5.1 |
| Proximal interphalangeal joint deformity | 4 | 2.9 |
| Bamboo spine | 3 | 2.2 |
| Arthritis mutilans | 3 | 2.2 |
| Others | 29 | 21.3 |
| Total | 136 | 100 |

 Table 5.11
 Factors associated with psoriatic arthritis

| Variable | | is Absent 5837) | Arthritis Present (n=1034) | | Multiple Logistic Regression ^a | | gression ^a |
|-----------------------------|------|--------------------|----------------------------------|--------|---|---------------|-----------------------|
| | n | (%) | N | (%) | Adj. OR | (95% CI) | P-value |
| Age: | | | | | | | < 0.001 |
| <18 years | 552 | (9.5) | 9 | (0.9) | 1.00 | - | - |
| 18-40 years | 2161 | (37.0) | 296 | (28.6) | 6.64 | (2.70, 16.35) | < 0.001 |
| 41-60 years | 2165 | (37.1) | 579 | (56.0) | 15.11 | (6.10, 37.47) | < 0.001 |
| >60 years | 959 | (16.4) | 150 | (14.5) | 11.80 | (4.60, 30.22) | < 0.001 |
| Age of onset: | | | | | | | |
| ≤40 years (Type 1) | 3850 | (66.6) | 707 | (68.8) | 1.39 | (1.10, 1.75) | 0.005 |
| >40 years (Type 2) | 1932 | (33.4) | 320 | (31.2) | 1.00 | - | - |
| Duration of disease: | | | | | | | |
| ≤5 years | 2779 | (48.1) | 300 | (29.2) | 1.00 | - | - |
| >5 years | 3003 | (51.9) | 727 | (70.8) | 1.54 | (1.26, 1.87) | < 0.001 |
| Gender: | | | | | | | |
| Male | 3371 | (57.8) | 516 | (49.9) | 1.00 | - | - |
| Female | 2466 | (42.2) | 518 | (50.1) | 1.65 | (1.40, 1.96) | < 0.001 |
| Ethnicity: | | | | | | | |
| Indian | 1020 | (17.5) | 227 | (22.0) | 1.30 | (1.06, 1.59) | 0.012 |
| Non-Indian | 4817 | (82.5) | 807 | (78.0) | 1.00 | - | - |
| Obesity group (WHO): | | | | | | | |
| BMI < 30 | 4468 | (80.7) | 735 | (74.6) | 1.00 | - | - |
| BMI≥30 | 1069 | (19.3) | 250 | (25.4) | 1.30 | (1.08, 1.57) | 0.007 |
| Type of psoriasis: | | | | | | | |
| Erythrodermic | 102 | (1.8) | 40 | (4.0) | | | NS |
| Non-erythrodermic | 5623 | (98.2) | 970 | (96.0) | | | |
| Body surface area: | | | | | | | |
| ≤10% | 3540 | (77.6) | 558 | (67.1) | 1.00 | - | - |
| >10% | 1019 | (22.4) | 274 | (32.9) | 1.24 | (1.02, 1.51) | 0.032 |
| Total skin score: | | | | | | | |
| <10 | 5398 | (93.4) | 907 | (89.0) | 1.00 | - | _ |
| ≥10 | 381 | (6.6) | 112 | (11.0) | 1.51 | (1.12, 2.03) | 0.006 |
| Nail involvement: | | | | | | | |
| Absence | 2519 | (43.5) | 221 | (21.8) | 1.00 | - | - |
| Presence | 3278 | (56.5) | 795 | (78.2) | 2.31 | (1.90, 2.81) | < 0.001 |
| DLQI: | | | | . , | | • | |
| ≤10 | 3742 | (66.4) | 590 | (58.4) | 1.00 | - | _ |
| >10 | 1897 | (33.6) | 420 | (41.6) | 1.48 | (1.24, 1.77) | < 0.001 |

^{*}Result was based on available information. Adj. OR = Adjusted odds ratio.^a Forward LR was applied. Multicollinearity was checked and not found. Hosmer-Lemeshow test(p=0.607), classification table (overall correctly classified percentage=84.6%) and area under the ROC curve (71.5%) were applied to check the model fitness.

CHAPTER 6

TREATMENTS

Types of treatment received by the patients for psoriasis in the last six months were analysed.

Most patients (96.3%) used some form of topical medications for psoriasis (**Table 6.1**). In 72.6% of the patients, topical monotherapy was the only treatment given. The most commonly used topical medication was topical steroids (85.4%). This was followed by topical tar preparation (80.7%), emollients (74.7%), keratolytics (54.8%) and vitamin D analogue such as calcipotriol (26.0%). Dithranol was less favoured and used in 1.9% of patients only (Table 6.2).

In the last six months prior to notification, 4.1% of patients received phototherapy (**Table** 6.3). Most of these patients (72.8%) were given narrowband UVB (NB-UVB) while 8.1% were given broadband UVB (BB-UVB). Less popular modalities were oral PUVA (4.0%), topical PUVA (1.1%) and bath PUVA (0.4%) (**Table 6.4**).

Systemic therapy was used in 19.6% of patients (Table 6.5). Methotrexate, being the commonest systemic therapy, was used in 67.8% of patients. This was followed by acitretin (20.8%), systemic corticosteroids (6.3%), sulphasalazine (6.2%), cyclosporine (4.5%), biologics (1.9%) and hydroxyurea (1.2%) (Table 6.6). 25 patients received biologic treatment. The biologic therapy most frequently used was etanercept (6 patients), followed by efalizumab and ustekinumab (4 patients each), adalimumab and infliximab (3 patients each). The name of the biologic agents was not specified in 5 patients.

Table 6.1 Use of topical therapy in psoriasis patients

| Topical therapy | No. | % |
|-----------------|------|-------|
| Yes | 6726 | 96.3 |
| No | 259 | 3.7 |
| Total | 6985 | 100.0 |

Distribution of types of topical therapy Table 6.2

| Topical therapy (one or multiple) | No. | % |
|---|-------|------|
| Topical steroids (other than face and flexures) | 5,742 | 85.4 |
| Tar preparation | 5,428 | 80.7 |
| Emollient | 5,021 | 74.7 |
| Keratolytics | 3,683 | 54.8 |
| Vitamin D analogues | 1,749 | 26.0 |
| Dithranol (anthralin) | 130 | 1.9 |
| Others | 160 | 2.4 |

Use of phototherapy in psoriasis patients **Table 6.3**

| Phototherapy | No. | % |
|--------------|--------|------|
| Yes | 272 | 4.1 |
| No | 6,435 | 95.9 |
| Total | 6, 707 | 100 |

Table 6.4 Distribution of types of phototherapy

| Type of phototherapy (one or multiple) | No. | % |
|--|-----|------|
| Narrowband UVB | 198 | 72.8 |
| Broadband UVB | 22 | 8.1 |
| Oral PUVA | 11 | 4.0 |
| Topical PUVA | 3 | 1.1 |
| Bath PUVA | 1 | 0.4 |
| Others | 2 | 0.7 |

Use of systemic therapy in psoriasis patients **Table 6.5**

| Systemic therapy | No. | % |
|------------------|--------|------|
| Yes | 1, 336 | 19.6 |
| No | 5, 493 | 80.4 |
| Total | 6, 829 | 100 |

Distribution of types of systemic therapy in psoriasis patients **Table 6.6**

| Type of systemic therapy (one or multiple) | No. | % |
|--|-----|------|
| Methotrexate | 906 | 67.8 |
| Acitretin | 278 | 20.8 |
| Sulphasalazine | 83 | 6.2 |
| Cyclosporin | 60 | 4.5 |
| Hydroxyurea | 16 | 1.2 |
| Biologics | 25 | 1.9 |
| Systemic corticosteroids | 84 | 6.3 |
| Others | 65 | 4.9 |

CHAPTER 7

QUALITY OF LIFE

There were a total of 4,295 adult patients (aged 17 and above) and 245 child/adolescent patients who completed the quality of life questionnaires, namely Dermatology Life Quality Index (DLQI) and Child Dermatology Life Quality Index (CDLQI).

The mean DLQI for adult psoriasis patients was 8.5 ± 6.4 , and the mean CDLQI for child/adolescent patients was 7.6 ± 5.5 .

The responses for each question of the DLOI and CDLOI were tabulated in **Table 7.1** and **7.2** respectively. 1,391 adult patients (32.4%) reported a DLQI of more than 10 indicating severe quality of life impairment due to psoriasis or its treatment (Figure 7.1). There were 230 adults (5.4%) who had a DLQI of more than 21 indicating extremely large effect on their quality of life by psoriasis. Nevertheless, 13.5% of adult patients reported no effect at all on their quality of life.

As shown in Figure 7.2, "systems and feelings" was the DLQI category most affected by psoriasis in adult patients. 37.5% of patients were affected very much or a lot by the itch and pain as well as embarrassment due to psoriasis. The aspect of life least affected by psoriasis was "personal relationship" in which 74.1% of the adult patients did not have or only have a little effect in this aspect.

In children/adolescents group, 18.0% of patients reported a CDLQI of more than 12 indicating very large or extremely large effect on QoL (Figure 7.3). There were 11 patients (4.5%) who had CDLQI of more than 19, reflecting extremely large effect of QoL. On the other hand, 12.7% child/adolescent patients reported no effect at all on their QoL.

In child/adolescent patients, the category of CDLQI most affected was "symptoms and feelings". 43.4% of children/adolescents reported that psoriasis affected very much or a lot in the symptoms and feelings domain. The aspect of life least affected by psoriasis was "personal relationship" in which 74.7% of the children did not have or only have a little effect (**Figure 7.4**). These results are similar to that of the adult patients.

Table 7.1 Responses for DLQI in adult psoriasis patients (age 17 and above)

| | | | | No. (%) | | |
|-----|--|--------------|-----------------|----------|------------|--------------|
| No. | DLQI Question | Very much | A lot | A little | Not at all | Not relevant |
| 1 | Over the last week, how itchy, sore, | 602 | 1,668 | 3,344 | 587 | 706 |
| | painful, or stinging has your skin been? | (9.5) | (26.4) | (52.9) | (11.1) | (11.2) |
| 2 | Over the last week, how embarrassed | 901 | 1,566 | 2,370 | 1,459 | |
| | or self conscious have you been | (14.3) | (24.9) | (37.6) | (23.2) | - |
| 2 | because of your skin? | , , | , , | , | , | |
| 3 | Over the last week, how much has your skin interfered with you going | 516 | 1 120 | 2,222 | 2,236 | 214 |
| | shopping or looking after your home | (8.2) | 1,138 (18.0) | (35.1) | | |
| | or garden? | (8.2) | (18.0) | (33.1) | (35.3) | (3.4) |
| 4 | Over the last week, how much has | 471 | 1,164 | 2,253 | 2,231 | 199 |
| | your skin influenced the clothes you | (7.5) | (18.4) | (35.7) | (35.3) | (3.1) |
| ~ | wear? | () | (==:) | (==::) | () | (=) |
| 5 | Over the last week, how much has | 554 | 1,177 | 2,158 | 2,220 | 208 |
| | your skin affected any social or leisure activities? | (8.8) | (18.6) | (34.2) | (35.1) | (3.3) |
| 6 | Over the last week, how much has | | | | | |
| U | your skin made it difficult for you to | 553 | 1,046 | 1,850 | 1,972 | 863 |
| | do any sport? | (8.8) | (16.6) | (29.4) | (31.4) | (13.7) |
| 7 | Over the last week, has your skin | | | | | |
| , | prevented you from working or | 536 | 405 | 1,426 | 2,341 | 923 |
| | studying? | (9.0) | (9.7) | (34.2) | (56.1) | (15.6) |
| 8 | Over the last week, how much has | | | | | |
| | your skin created problems with your | 273 | 710 | 1,998 | 3,050 | 279 |
| | partner or any of your close friends or | (4.3) | (11.3) | (31.7) | (48.3) | (4.4) |
| | relatives? | , , | ` , | , , | , , | ` ' |
| 9 | Over the last week, how much has | 203 | 329 | 1,113 | 3,078 | 1,530 |
| | your skin caused sexual difficulties? | (3.2) | (5.3) | (17.8) | (49.2) | (24.5) |
| 10 | Over the last week, how much of a | | | | | |
| | problem has the treatment for your | 339 | 965 | 2,295 | 2,378 | 334 |
| | skin been, for example by making | (5.4) | (15.3) | (36.4) | (37.7) | (5.3) |
| | your home messy or by taking up time? | (3.4) | (13.3) | (30.4) | (31.1) | (3.3) |




Figure 7.1 Quality of life in adult patients with psoriasis




Figure 7.2 QoL impairment in adults psoriasis patients based on category of DLQI

Responses for CDLQI in child/adolescent psoriasis patients (aged 5 to 16) **Table 7.2**

| | | | | No. (%) | | |
|-----|---|--------------|---------------|---------------|---------------|--------------|
| No. | CDLQI Question | Very much | A lot | A little | Not at all | Not relevant |
| 1 | Over the last week, how itchy, "scratchy", sore, painful, or stinging has your skin been? | 29 (7.3) | 116 (29.4) | 215 (54.4) | 35 (8.9) | - |
| 2 | Over the last week, how embarrassed or self conscious have you been because of your skin? | 61 (15.5) | 101 (25.6) | 158 (40.1) | 74 (18.8) | - |
| 3 | Over the last week, how much has your skin affected your friendships? Over the last week, how much have | 16 (4.1) | 48 (12.2) | 133 (33.8) | 197 (50.0) | - |
| 4 | you changed or worn different or special clothes/shoes because of your skin? | 18 (4.6) | 60 (15.3) | 144 (36.6) | 171 (43.5) | - |
| 5 | Over the last week, how much has your skin trouble affected going out, playing, or doing hobbies? | 20 (5.1) | 63 (16.0) | 146 (37.1) | 165 (41.9) | - |
| 6 | Over the last week, how much have you avoided swimming or other sports because of your skin trouble? | 24 (6.1) | 56 (14.2) | 115 (29.3) | 198 (50.4) | - |
| 7 | If school time: Over the last week, how much did your skin problem affect your school work? Or If holiday time: Over the last week, has your skin problem interfered with your enjoyment of the holiday? Over the last week, how much trouble | 14 (3.6) | 48 (12.3) | 133 (34.0) | 196 (50.1) | - |
| Ü | have you had because of your skin with other people calling you names, teasing, bullying, asking questions or avoiding you? | 17 (4.3) | 36 (9.2) | 101 (25.7) | 239 (60.8) | - |
| 9 | Over the last week, how much has your sleep been affected by your skin problem? | 24 (6.4) | 46 (12.3) | 128 (34.2) | 176 (47.1) | - |
| 10 | Over the last week, how much of a problem has the treatment for your skin been? | 16 (4.1) | 52 (13.3) | 143 (36.6) | 180 (46.0) | - |




Figure 7.3 Quality of life in children/adolescents with psoriasis.




Figure 7.4 QoL impairment in child/adolescent psoriasis patients based on category of DLQI

Chapter 8: Outcomes

CHAPTER 8

OUTCOMES

In this registry, follow-up data were collected approximately every 6 months. Outcomes of patients were assessed by measuring the change in several clinical parameters between the last follow-up visit and the visit at registration. Severity of psoriasis skin lesions were assessed in terms of the extent of lesions, i.e. percentage of body surface area involvement, and lesional characteristics via clinical skin scoring method for each of the five body regions. Other clinical parameters monitored include severity of joint pain on a visual analogue score (0-10), and quality of life using Dermatology Life Quality Index (DLQI).

From a total of 6,985 psoriasis patients registered in MPR, follow-up data were obtained in 1,936 patients. The mean duration of follow-up was 19.7 ± 12.2 months, with the longest duration of 36 months (**Table 8.1**).

Extent of Psoriasis Lesions

The extent of psoriasis lesions was assessed in terms of percentage of body surface area involvement categorised into 4 scales, i.e. <5%, 5%-10%, 10%-90%, and >90% (erythrodermic). A total of 1, 342 patients were evaluated for change in the extent of lesions. Of these patients, 318 patients (23.7%) had improvement by at least one scale, among which 57 (4.2%) had improvement by two scales, and 3 patients improved from BSA>90% to BSA<2%. No improvement was found in 721 patients (53.7%), and 243 patients (18.1%) had worsening by at least one scale (**Figure 8.1**).

Clinical Skin Scores

Clinical skin scores measures the thickness, erythema and scaliness of the psoriasis lesions in each of the five body regions. A score of 0 to 3 is given for each body region. Total Clinical Skin Score is the total of the scores in all five body regions. 124 patients (7.7%) had the most marked improvement in skin scores by 75% or more, and 364 patients (22.6%) had improvement by 50-75%, while 335 patients (14.9%) had 25-50% improvement. 166 patients (11.1%) had modest improvement of less than 25%. No improvement of skin scores were detected in 359 patients (19.5%). Skin scores worsened in 535 patients (24.1%) (**Figure 8.2**).

Joint Pain

From a total of 130 patients who reported to have joint pain, 63 patients (48.5%) had improvement in joint pain as measured by the visual analogue scale. Of these patients, 21 patients (16.2%) had improvement of between 50% and 75%, 2 patients (1.5%) had improvement of more than 75%, 27 patients (20.8%) had improvement of between 25% and 50%, and 13 patients (10.0%) had improvement of less than 25%. There was no improvement of joint pain in 31 patients (23.8%), while joint pain worsened in 36 patients (27.7%) (**Figure 8.3**).

Change in Quality of Life

In adult patients aged 17 years and above, we noted an overall improvement in the quality of life. A total of 1,791 adult patients were evaluated for change in quality of life by DLQI. Of these patients, 412 patients (23.0%) had significant improvement with a reduction of DLQI score by at least 5, whereas 289 patients (16.1%) had significant worsening with an increase in DLQI score by at least 5 (**Figure 8.4**).

A total of 61 patients aged below 17 were evaluated for change in quality of life by DLQI. Of these patients, 14 patients (23.0%) had a significant improvement of Child DLQI score by at least 5, while 5 patients (8.3%) worsened (Figure 8.4).

Table 8.1 Distribution of psoriasis patients according to the duration of follow-up

| Duration of follow-up | No. | % |
|------------------------------|------|-------|
| 0 to 6 months | 271 | 14.0 |
| 7 to 12 months | 459 | 23.7 |
| 13 to 18 months | 362 | 18.7 |
| 19 to 24 months | 267 | 13.8 |
| 25 to 30 months | 192 | 9.9 |
| 31 to 36 | 162 | 7.9 |
| >36 | 233 | 12.0 |
| | 1936 | 100.0 |

Mean duration of follow-up: 15.5 ± 8.4 months (range 0 - 36 months)




Figure 8.1 Improvement in the extent of skin lesions




Figure 8.2 Improvement in the total clinical skin scores




Figure 8.3 Improvement in joint pain




Figure 8.4 Improvement in DLQI and CDLQI

Chapter 8: Outcomes

APPENDIX A: CASE REPORT FORM

Case Report Form

| Instruction: Where o | NATIONAL DERMATOLOGY REGISTRY (Derm Reg) Malaysian Psoriasis Registry Case Report Form Instruction: Where check boxes № 1 are provided, check (¬1) one or more boxes. Where radio buttons are provided, check (¬1) one button only. | | | | | | | | | | | |
|--|--|---|--|--|--|--|--|--|--|--|--|--|
| Doctor's Name : | | | | | | | | | | | | |
| Name of Institution : | | | | | | | | | | | | |
| SECTION 1: DE | MOGRAPHIC DETAILS | | | | | | | | | | | |
| Patient visit date: (dd/mm/yyyy) | 2. Type of visit : | New Case Follow-Up | | | | | | | | | | |
| 3. Name of patient: | | | | | | | | | | | | |
| 4. NRIC : | MyKad/ MyKid: | Old IC: | | | | | | | | | | |
| | type (if others): Opassport Father's I | pecify document | | | | | | | | | | |
| 5. Address: | Town / City: St | ate: | | | | | | | | | | |
| 6. Contact #number: | Homephone: - | H/P: - | | | | | | | | | | |
| 7. Gender: # | ◯ Male ◯ Female | | | | | | | | | | | |
| 8. Date of birth : # (dd/mm/yyyy) | / / Estimated/ presume | dyear If the exact date is not known, please enter 01/67/3539 & check the estimated presumed year tox | | | | | | | | | | |
| 9. Ethnic group : ∉ | ◯ Malay ◯ Chinese ◯ Indian ◯ | O Malay O Chinese O Indian O Crang Asli O Others, specify: | | | | | | | | | | |
| 10. Nationality : | ◯ Malaysian ◯ Non-Malaysian, specify | | | | | | | | | | | |
| 11. Marital status : # | ◯ Single ◯ Married ◯ Divorced ◯ | Widow Widower | | | | | | | | | | |
| SECTION 2 : ME | DICAL HISTORY | | | | | | | | | | | |
| 1. Age when # psoriasis started: | 2. Age wher # psoriasis diagnose | | | | | | | | | | | |
| 3. Family member(s) with psoriasis: | ○ No ○ Yes → ☐ Father ☐ Siblin (if YES, please 6ck ONE or MULTIPLE) ☐ Mother ☐ Childs | | | | | | | | | | | |
| Aggravating factors: | No | | | | | | | | | | | |
| 5. Disease burden in the | a) No.of clinic visits due to psoriasis : | (enter 0 if none) | | | | | | | | | | |
| last 6 months : | b) No. of days off work / school due to psoriasis : | (enter 0 if none) Not applicable | | | | | | | | | | |
| 6. Other | c) No. of hospital admissions due to psoriasis : a) Ischaemic heart disease : | (enter Offmone) | | | | | | | | | | |
| diseases: | a) Ischaemic neart disease : b) Cerebrovascular disease (stroke) : | Yes No Unknown | | | | | | | | | | |
| | c) Diabetes mellitus : | Yes No Unknown | | | | | | | | | | |
| | d) Hypertension : | Yes No Unknown | | | | | | | | | | |
| | e) Hyperlipidaemia : | Yes No Unknown Yes No Unknown | | | | | | | | | | |
| | f) Other diseases, specify: (e.g. HV infection, tuberculosis, lymphoma, etc.) | Yes No Unknown | | | | | | | | | | |
| 7. Cigarette smoking : | ○ Never smoked ○ Ex-smoker ○ Current smoker | cigarettes per day | | | | | | | | | | |

Items marked # above need not be entered if the patient has been previously notified to the registry

Version 2.4 Last updated 07/12/2011

| N | NATIONAL DERMATOLOGY REGISTRY (DermReg) Malaysian Psoriasis Registry Case Report Form | | | | | | | | | | | |
|--|---|---------------------------------|------------------------|---------------------|---|----------------------------------|----------------|-----------------------|--|--|--|--|
| Instruction: Where check boxes 1 are provided, check (1) one or more boxes. Where radio buttons are provided, check (1) one button only. | | | | | | | | | | | | |
| | | | | | 1 | | | | | | | |
| | INICAL EXAMINA | ATION | | lor s s | | | | | | | | |
| 1. (a) Height: | (cm) | | | (b) \ | Weight: | | (kg) | | | | | |
| 2. Type of psoriasis: | (Please select ONE pred Plaque Generalised pu | ⊕ Guttate ustular ⊜ Localise | _ | ythroderm | nic ⊜ Flexural/ ⊝ Palmopla | | _ | thers,specify: | | | | |
| 3. Severity: | Body surface area i | nvolved: 🔘 < | <5% 🔘 5 | - 10% | ○ > 10% | 0 |) Erythrode | rmic (>90%) | | | | |
| | Body part | Grade o | of severity | | Key for grading | | | | | | | |
| | Scalp | ○ ○ ○ 1 | ② 2 ③ 3 | 1 | Grade 0 : Skin normal or hypo-/hyperpigmented patch only. | | | | | | | |
| | Face & Neck | ○ 0 0 1 | | ı | | erythema,fine s ral clearing. | cales, thin pl | aque, with or without | | | | |
| | Trunk | ○ 0 ○ 1 | | ı | Grade 2 : Mod | erale erythema | or scaling, n | moderately thick | | | | |
| | Upper Limbs | ○ 0 0 1 | | | plaq Grade 3 : Seve | ue. ere erythema or | scaling, ven | v thick plaque | | | | |
| | Lower Limbs | ○ 0 ○ 1 | | ı | | , | | , | | | | |
| 4. Nail involvement : | ○ No ○ `` (ii YES, please tick ONE | , , | Pitting Onycholysis | ☐ Discol ☐ Subur | loration ngual hyperkerate | | tal nail dys | trophy | | | | |
| 5. Joint | ○ No ○ ' | Yes → | | | | | | | | | | |
| disease : | a) Rheumatoid facto | O regulate | | Positive | ve ⊚ Not/ | Available | | | | | | |
| | b) Morning stiffness | > 30 minutes | | | ○ No | 0 1 | es es | | | | | |
| | c) Enthesopathy / Da | 'es | | | | | | | | | | |
| | d) Type :- | 1. Oligo√ Mon | oarthropathy | | ○ No | () Y | 'es | | | | | |
| | | 2. Distal hand | | | ○ No | () | 'es | | | | | |
| | | 3. Symmetrical (Rheumatoid | | thy | ○ No | 01 | es | | | | | |
| | | 4. Spondylitis/ | <u> </u> | | ○ No | 0 1 | /es | | | | | |
| | | 5. Arthritis mut | tilans | | ○ No | ○ No ○ Yes | | | | | | |
| | e) Severity:- | 1. Pain | () No (|) Yes - | | ore (1-10): | | | | | | |
| | | 2. Swelling | - | | - | | | | | | | |
| | | | ⊕ N0 (| ⊚ Yes | Please Specify: | | | | | | | |
| | | 3. Deformity | ⊚ No (| Yes - | → Please Sp | | | | | | | |
| SECTION 4 : TI | REATMENT REC | EIVED IN THE F | PAST 6 MO | NTHS | | | | | | | | |
| 1. Topical | a) Tar preparation | ⊕ No | ○ Yes | | al steroids | | (No | | | | | |
| therapy: | b) Vitamin D analog | In a | | _ | than face / flexures; olytics e.g. salic | | - | () Yes | | | | |
| | e.g calcipotriol | ○ No | ○ Yes | g) Emol | | yne dele | ○ No | ① Yes | | | | |
| | c) Calcipotriol with betamethasone | ○ No | Yes | - | rs, specify | | - | | | | | |
| | dipropionate | | | in ourer | o, opeony | | ⊚ No | ⊚ Yes 🔻 | | | | |
| | d) Dithranol (anthra | llin) 🔘 No | Yes | 1 | | | <u> </u> | | | | | |
| 2. Phototherapy : | ○ No | ○ Yes → | ☐ BB-UVB | | oral PUVA | Topical PUVA | Other | s,specify | | | | |
| | (if YES, please fick O | NE or MULTIPLE) | ■ NB-UVB | _ | | Excimer lase | | | | | | |
| 3. Systemic | ○ No | Yes | <u>'</u> | | | | | | | | | |
| therapy: | a) Methotrexate | ⊝ No | Yes | f) Biolog | gics, specify | | ○ No | ⊚ Yes 🕞 | | | | |
| | b) Acitretin | ○ No | ○ Yes | | | | [| | | | | |
| | c) Sulphasalazine | ○ No | ○ Yes | g) Syste | emic corticoster | oids | ○ No ○ Yes | | | | | |
| | d) Cyclosporin | ○ No | Yes | - | rs, specify | | ○ No ○ Yes → | | | | | |
| | e) Hydroxyurea | ○ No | Yes | , , , , , , , | , -, , | | i | <u> </u> | | | | |
| SECTION 5: OF | IALITY OF LIFE | 1 | | | | | <u></u> | | | | | |
| 1. Quality of Life : | Diseas instruct | and assist actions in | completion th | a attacke | d DLOI form | | | | | | | |
| i. Guarry of Life : | Please Instruct | and assist patient in | completing th | e attache | O DEGI IOIM | | | | | | | |

**Note: Please ensure that all sections of this form have been completed. Kindly submit to: Malaysian Pscriasis Registry, Department of Dermatology, Hospital Kuala Lumpur, Jalan Pahang, 50586 Kuala Lumpur

Version 2.4 Last updated 07/12/2011

NATIONAL DERMATOLOGY REGISTRY (DermReg) CONFIDENTIAL Malaysian Psoriasis Registry Dermatology Life Quality Index (DLQI) (For Adults of Age 17 and Above) Instruction: Where check boxes \bigvee are provided, check $(\sqrt[4])$ one or more boxes. Where radio buttons \bigcirc are provided, check $(\sqrt[4])$ one button only.

Objektif kaji ælidik adalah untuk memahami setakat manakah masalah kulit anda mempengaruhi kehidupan anda SEPANJANG MINGGU LALU. The aim of this questionnaire is to measure how much your skin problem has affected your life OVER THE LAST WEEK.

| Sila tandakan satu kotak (i) untuk ætiap soalan / Please tick $^{\alpha} \sqrt{^{\alpha}}$ one box for each qu | estion 清在每个 | 何題后选择 | 一项打"√". | DLQI Score | Auto calculated |
|--|--------------------------------------|----------------------|---------------------------|--|---|
| Sepanjang Minggu Lalu OVER THE LAST WEEK 上周内, | Sangat Banyak Very much 非常多 | Banyak Alot 许多 | Sedikit A little 一点 | Tidak Langsung <i>Not at all</i> 完全没有 | Tidak Berkenaan Not Relevant 无关 |
| Setakat manakah kulit anda berasa gatal atau sakit? Over the last week, how itchy, sore, painful or stinging has your skin been? 您的皮肤感到痒、触痛、疼痛、刺痛了吗。? | 0 | 0 | 0 | 0 | |
| 2) Setakat manakah anda berasa malu atau segan, disebabkan oleh kulit anda? Over the last week, how embarrassed or self conscious have you been because of your skin? | 0 | 0 | 0 | 0 | |
| 由于您的皮肤问题,您感到尴尬或自卑吗? | | | | | |
| 3) Setakat manakah kulit anda menganggu anda daripada pergi membeli belah atau menjaga rumah atau berkebun? Over the last week, how much has your skin interfered with you going shopping or looking after your home or garden? | 0 | 0 | 0 | 0 | 0 |
| 因为皮肤问题、对您购物、做家务、整理庭院影响程度如何? 4 Setakat manakah kulit anda mempengaruhi pakaian yang anda pakai? | | | | | |
| Over the last week, how much has your skin influenced the clothes you wear? 皮肤问题对您穿衣服影响程度如何? | 0 | 0 | 0 | 0 | 0 |
| 5) Setakat manakah kulit anda mengganggu aktiviti - aktiviti sosial atau masa lapang anda ? | | | | | |
| Over the last week, how much has your skin affected any social or leisure activities? | 0 | 0 | 0 | 0 | 0 |
| 皮肤问题对您的社交或休闲生活有多大的影响? 6) Setakat manakah keadaan kulit anda menyebabkan anda tidak selesa | | | | | |
| bersukan? Over the last week, how much has your skin made it difficult for you to do any sport? 皮肤问题对您运动有多大妨碍? | 0 | 0 | • | 0 | 0 |
| 7) Adakah kulit anda menyebabkan anda tidak bekerja atau belajar? | | | | | |
| Over the last week, has your skin prevented you from working or studying? | | | | | |
| 皮肤问题是否让您无法上班或学习? | | | | | |
| ■ YaYes是 ■ Tidak No 不是 ■ Tidak Berkenaan Not Relevant 无关 | | | | | |
| *Jika *tidak*, setakat manakah kulit anda menjadi masalah semasa kerja atau belajar? | | ļ | | ļ | |
| If "No", over the last week how much has your skin been a problem at work or studying? | | 0 | 0 | 0 | |
| 如果选择 "不是",那么上周内您的皮肤问题对工作或 学习有 多大影响呢? | | | | <u> </u> | |
| 8) Setakat manakah kulit anda menimbulkan masalah dengan teman, rakan baik atau saudara mara anda? | | | | | |
| Over the last week, how much has your skin created problems with your partner or any of your close friends or relatives ? | • | 0 | 0 | 0 | 0 |
| 皮肤问题妨碍了您和爱人、亲密的朋友、亲戚间的交往了 吗? 9) Setakat manakah kulit anda menyebabkan sebarang masalah hubungan seks? Over the last week, how much has your skin caused sexual difficulties? | 0 | 0 | 0 | 0 | 0 |
| 皮肤问题给您的性生活造成了多大影响? | | | | | |
| 10) Setakat manakah rawatan kulit anda menimbulkan masalah seperti mengotori rumah anda atau mengambil masa anda? | | | | | |
| Over the last week, how much of a problem has the treatment for your skin been, for example by making your home messy or by taking up time? 由于治疗您皮肤的毛统。给您造成了多少麻烦、如把家 里弄得一团 | 0 | 0 | 0 | 0 | 0 |

Sila semak sama ada SETIAP soalan telah dijawab. Terima kasih Please check you have answered EVERY question. Thank you.

请您检查您是否已回答所有问题. 谢谢合作

Version 2.4 Last updated 07/12/2011 @AY Finlay, GK Khan, April 1992, This must not be copied without the permission of the authors. page 3 of 4

NATIONAL DERMATOLOGY REGISTRY (DermReg) CONFIDENTIAL For Office Use only: Malaysian Psoriasis Registry Children's Dermatology Life Quality Index (DLQI) (For age 5 to 16) Instruction: Where check boxes M are provided, check (\(\sqrt{)}\) one or more boxes. Where radio buttons \(\infty\) are provided, check (\(\sqrt{)}\) one button only.

Objektif kaji ælidik adalah untuk memahami setakat manakah masalah kulit anda mempengaruhi kehidupan anda SEPANJANG MINGGU LALU. The aim of this questionnaire is to measure how much your skin problem has affected your life OVER THE LAST WEEK.

| 这份问卷的目的是衡量上周内您的皮肤问题对您的生活造成了多大的影响。 | | DI | Ol Cassas | Auto |
|--|---------------------|-----------------------|------------|------------------------|
| Sila tandakan satu kotak (√) untuk setiap soalan / Please tick "√" one box for each question清在每 | 个问题后选择一 | -项打"√" <mark>□</mark> | .QI Score: | calculated |
| Sepanjang Minggu Lalu | Sangat | Banyak | Sedikit | Tidak |
| OVER THE LAST WEEK 过去一星期中 | Banyak Very much | A lot | A little | Langsung Not at all |
| 双女一生用于 | 非常多 | 许多 | 一点 | 完全没有 |
| 1) Setakat manakah kulit anda berasa gatal atau sakit ? | 11-70-37 | 112 | | 70±10,11 |
| Over the last week, how itchy, "scratchy", sore or painful has your skin been? | 0 | | 0 | 0 |
| 你皮肤发痒、搔抓、破皮或疼痛的程度是如何? | | | | |
| 2) Setakat manakah anda berasa malu, segan, susah hati atau sedih disebabkan | | | | |
| oleh kulit anda? | | | | |
| Over the last week, how embarrassed or self conscious, upset or sad have you been because of your skin? | 0 | 0 | 0 | 0 |
| 你因为自己皮肤问题而感到难为情或害羞、苦恼或难过的程度是如何? | _ | | | |
| Setakat manakah kulit anda mempengaruhi persahabatan anda? | | | | |
| Over the last week, how much has your skin affected your friendships? | 0 | 0 | 0 | 0 |
| 皮肤问题对你和朋友交往的影响是如何? | | | | |
| 4) Setakat manakah anda menukar atau memakai pakaian atau kasut kerana kulit anda? | | | | |
| Over the last week, how much have you changed or worn different or special | 0 | | | |
| clothes/shoes because of your skin? | | 0 | 0 | 0 |
| 你因为皮肤问题而改变穿著不同或特定衣鞋的影响是如何? | | | | |
| Setakat manakah masalah kulit anda mempengaruhi anda untuk keluar, bermain atau melakukan hobi anda? | | | | |
| | _ | _ | | |
| Over the last week, how much has your skin trouble affected going out playing, or doing hobbies? | 0 | 0 | 0 | 0 |
| 皮肤的问题对你外出、玩耍、或从事休闲嗜好的影响是如何? | | | | |
| 6) Setakat manakah anda menjauhi diri daripada berenang atau melakukan sukan lain | | | | |
| disebabkan oleh masalah kulit anda? | | | | |
| Over the last week, how much have you avoided swimming or other sports because of your skin trouble? | | | | 0 |
| 你因为皮肤的问题而避免游泳或其他运动的影响程度是如何? | | _ | | _ |
| 7).Pada minggu yang lalu, | | | | |
| Last week, 过去一星期 | | | | |
| Pada hari persekolahan, setakat manakah kulit anda mempengaruhi kerja sekolah anda? | | | | |
| If school time: Over the last week, how much did your skin problem affect your school | | | | |
| work? | | | | |
| 如果是上课时间,皮肤问题影响你学校功课的程度是如何? ATAU OR 並 | 0 | 0 | | 0 |
| Pada hari cuti, setakat manekah kulit anda mengganggu anda menikmati cuti? | | | | |
| If holiday time: Over the last week, has your skin problem interfered with your | | | | |
| enjoyment of the holiday? | | | | |
| 如果是放假期间,皮肤问题干扰到你享受假期的兴致是如何? | | | | |
| 8) Setakat manakah orang menggelar anda dengan nama yang tidak baik, mengejek, | | | | |
| menanya soalan-soalan atau menjauhi diri disebabkan oleh kulit anda? | | | | |
| Over the last week, how much trouble have you had because of your skin with other | | 0 | | 0 |
| people calling you names, teasing, bullying, asking questions or avoiding you? | _ | | | |
| 因为皮肤的问题使得别人骂你、嘲笑你、欺负你、问你问题或躲避你,这种困扰程度是如何? 9) Setakat manakah masa tidur anda diganggu kerana masalah kulit? | | | | |
| Over the last week, how much has your sleep been affected by your skin problem? | | | _ | _ |
| 你因皮肤的问题而影响到睡眠的程度是如何? | 0 | 0 | 0 | 0 |
| 10) Setakat manakah rawatan kulit anda menjadi suatu masalah? | | | | |
| Over the last week, how much of a problem has the treatment for your skin been? | 0 | | | |
| 针对皮肤所进行的治疗对你产生的困扰程度是如何? | | 0 | 9 | |
| The second section of the second seco | | | | |

Sila semak sama ada SETIAP soalan telah dijawab. Terima kasih Please check you have answered EVERY question. Thank you. 请您检查您是否已回答所有问题。谢谢合作

GM.S. Lewis-Jones, A.Y. Finlay, May 1993, This must not be copied without the permission of the authors. page 4 of 4

Biologic Treatment Initiation Form

| NATIONAL D | ERMATO | LOGY REGISTRY | (DermReg) | | CONFIDE | NTIAL | | | | |
|--|----------------|--|-----------------------|---|---------------|------------------|--|--|--|--|
| M | alaysian | Psoriasis Registry | | | al use only: | | | | | |
| | | tment Initiation Fo | | ID: | | | | | | |
| Instructions: Where check b radio buttons | | orovided, check (✓)one o des, check (✓) one butto | | Centre | | | | | | |
| | | | | | | | | | | |
| Doctors' name | | | | | | | | | | |
| Name of institution | | | | | | | | | | |
| | | | | | | | | | | |
| DEMOGRAPHIC DE | TAILS | | | | | | | | | |
| 1. Visit date (dd/mm/yyyy) | | / | | | | | | | | |
| 2. Name of patient | | | | | | | | | | |
| | MyKad/ MyP | Gd: | | C | Nd IC: | | | | | |
| | Other ID do | cument No: | | | | | | | | |
| 3. NRIC | Specify doc | ument @ Registration nun | nber @ Mother's I/C | Work Per | mit @ Clin | ric RN | | | | |
| | type (if other | | | Driver's Li | cence @ Poli | ice ID Card | | | | |
| | | Birth Certificate | Armed Force I | ID Hospital F | IN @ Oth | ers | | | | |
| | | | | | | | | | | |
| SECTION 1 : MEDIC | AL HISTO | RY | | | | | | | | |
| 1. Any history of tubercu | ılosis? | ⊚ No ⊚ Yes | If yes, please speci | fy: | | | | | | |
| | | Date of diagnosis (dd/mm/yyyy) | | | | | | | | |
| | | Type of tuberculosis | | | 1 | | | | | |
| | | - | Latent tuberor | | | | | | | |
| | | | Pulmonary tul | | | | | | | |
| | | | | ary tuberculosis | | | | | | |
| | | | Specify organ | n involved: | | | | | | |
| | | Completed anti-TB treatment? | @ No | /es | | | | | | |
| 2. Any history of cancer | ? | ® No ® Yes, sp | ecify | | | | | | | |
| 3. Any history of neurolo disease? | gical | ® No ® Yes, sp | ecify | | | | | | | |
| 4. Any history of liver dis | | ⊚ No ⊗ Yes, sp | ecify | | | | | | | |
| 5. Any history of cardiov disease? | ascular | ⊚ No ⊗ Yes, sp | ecify | | | | | | | |
| 6. Previous systemic tre | atment? | ⊚ No ⊚ Yes I | f yes, please specify | | | | | | | |
| | | Systemic a | agent | Re | asons for sto | pping | | | | |
| | | ☐ Phototherapy | | ☐Poor response ☐Others (specify) _ | □intolerance | ☐Adverse effects | | | | |
| | | Oral methotrexate | | ☐Poor response ☐Others (specify) | □intolerance | Adverse effects | | | | |
| | L | ☐ Parenteral methotrex | ate | ☐Poor response ☐Others (specify) | □intolerance | Adverse effects | | | | |
| | ⊢ | □ Acitretin | | ☐Poor response ☐Others (specify) ☐Poor response | □Intolerance | ☐Adverse effects | | | | |
| | <u></u> | □ Sulphasalazine | | Cothers (specify) | Dintolerance | DAdverse effects | | | | |
| | - | Cyclosporin | | □Others (specify) _ □Poor response | □intolerance | ☐Adverse effects | | | | |
| | | ☐ Hydroxyurea ☐ Systemic corticostero | ide | □Others (specify) □Poor response | □intolerance | ☐Adverse effects | | | | |
| | - ⊢ | ☐ Biologic therapy, spec | | ☐Others (specify) ☐ ☐Poor response | □intolerance | ©Adverse effects | | | | |
| | - ⊢ | Other systemic agent | , | □Others (specify) □ □Poor response □Others (specify) | □intolerance | GAdverse effects | | | | |

| SECTION 2 : PRE-TREATMENT PASI EVALUATION | | | | | | | | | | | | | | | | | | |
|---|----------|------|--------|--------|----------|----------------------------------|--------|----------|-------|---------|---------|-------|--------|-------|--------|---------------------------|----------------------------|--|
| SECTION 2:11 | 1 | \L, | | | | | | | | eristic | | | | | | 1 | | |
| Body region | | 0 | - 14 | one. | 4-1 | | | | | | re, 4 = | Von | reau | oro | | | nvolvement of | |
| Dody region | \vdash | | | ema | 1 - N | 10, 2 - | | ickn | | | re, 4 - | | calir | | | each bo | each body region | |
| | ┼ | EI | yune | ama | | ├ | - " | 255 | | _ | - | Calli | 'y | | ⊙ None | o 50 − 69% | | |
| Head | 0 | 0 | 0 | 0 | 0 | | 0 | | 0 | | | | 0 | | | @ 1 - 9% | © 70 – 89% | |
| neau | 0 | 1 | 2 | 3 | 4 | 0 | 1 | 2 | 3 | 4 | 0 | 1 | 2 | 3 | 4 | © 10 - 29% | 90 – 100% | |
| | +- | | | | | \vdash | | | | | | | | | | ⊙ 30 – 49% ⊙ None | ⊚ 50 − 69% | |
| Upper limbs | 0 | 0 | 0 | ⊚ 3 | 0 | 0 | ⊚ 1 | 0 | 0 | • | 0 | 0 | ⊚ 2 | 0 | 0 | ⊚ 1 - 9% | ⊙ 70 – 89% | |
| opper minos | 0 | 1 | 2 | 3 | 4 | 0 | 1 | 2 | 3 | 4 | 0 | 1 | 2 | 3 | 4 | ⊙ 10 - 29% ⊙ 30 - 49% | ⊚ 90 – 100% | |
| | | | | | | <u> </u> | | | | _ | | | | | _ | None | ⊙ 50 − 69% | |
| Trunk | | | | 3 | | | ⊚ 1 | | | | | | 2 | | | ⊚ 1 - 9% ⊚ 10 - 29% | ⊙ 70 – 89% ⊙ 90 – 100% | |
| | Ľ | | _ | | | Ĭ | | _ | _ | | ľ | | _ | _ | | 30 − 49% | | |
| | | | | 0 | | | | | _ | • | | _ | 0 | | | None 1 - 9% | ⊚ 50 – 69% ⊚ 70 – 89% | |
| Lower limbs | l ő | | | 3 | | | 1 | | | | | | 2 | | | ⊚ 10 - 29% | | |
| | | | | | | | | | | | | | | | | @ 30 - 49% | | |
| | | | | | | | | | | | | | | | | | | |
| SECTION 3 : IN | VES | TIG. | ATI | ON | <u> </u> | | | | | | | | | | | | | |
| 1. Mantoux test | | | | _ | | | mm | | Not (| done b | ecause | of_ | | | | | | |
| 2. Interferon-γ rele | ase a | ssay | | @ N | legati | ative ⊚ Positive ⊚ Not available | | | | | | | | | | | | |
| | Norr | | | | lorma | | | ⊚ AŁ | non | mal | ΘN | ot a | vaila | ble | | | | |
| 3. Chest X-ray | | | | lf al | nom | al, spe | cify f | indin | ms | | | | | | | | | |
| | | | | II CII | JI KATI | ai, spe | Oily I | III CIII | 95_ | | | | | | | | | |
| 4. Hepatitis B stat | us | | | @ N | legati | ve | (| 9 Po | sitiv | e | ⊚ N | ot a | vaila | ble | | | | |
| 5. Hepatitis C stat | us | | \Box | Θľ | legati | tive | | | | | | | | | | | | |
| 6. HIV status | | | | @ N | legati | ative © Positive © Not available | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | |
| SECTION 4 : IN Indication for biol | | | | | BIC | LOG | | | | | | | | | | y are contraindic | nt nd | |
| indication for bior | ogic u | eau | leni | | - | - | | | | | | | _ | | | temic therapy | atieu | |
| | | | | | - | - | | | | | | | | | _ | | | |
| | | | | | - | - | _ | | _ | | | Star | luaro | ı sys | aemic | therapy | | |
| | | | | | | _ | I۲ | uier | indic | ation, | specify | | | | | | | |
| | | | | | | | ╚ | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | |
| SECTION 5 : C | URRI | ENT | BI | OLO | GIC | TRE | ATN | 1EN | T | | | | | | | | | |
| 1. Biologic agent | | | | | | ⊚ Infli | dmal | ١V | | | 0 | Eta | nero | ept S | SC. | | | |
| | | | | | | | | nab | SC | | 0 | Ust | ekin | umal | b SC | | | |
| | | | | | | ⊚ Othe | ers. s | peci | fv | | | | | | | | | |
| 2. Date start (dd/mm/yyyy) | | | | | | |]/[| |]/[| | |] | | | | | | |
| 3. Dose | | | | | | | | | | | | | | | | | | |
| <u> </u> | | | | | | | | | | | | | | | | | | |
| SECTION 6 : CONCOMITANT SYSTEMIC TREATMENT | | | | | | | | | | | | | | | | | | |
| Concomitant systemic treatment / phototherapy | | | | | | ⊚ No | | 0 | Yes | s, spec | ify | | | | | | | |
| | | | | | | | | | | | | | | | | | | |

Biologic Treatment Follow Up Form

| NATION | | | | |)LOG Psori | | | | | ermF | leg) | | | For | official | CONFIDENTIAL al use only: | | |
|----------------------------------|---------------|--------------|------------------|-------------|----------------------|--------------------|------------|---------------|--------------|-----------------|-----------------------|--------|----------------------|--------|-------------------------|---------------------------------------|----------------------------|--|
| | Biol | ogi | c Tr | reat | ment | Follo | wŪ | рF | оп | | | | | ID: | | | | |
| Instructions: Where c radio b | heck utton | baxa ıs ⊙ | es 🗆 are p | are rovi | provideo des, che | i, check ck (√) | (√) one | lone butti | or m m or | iore bo: nly | ces. Wh | ете | | Cer | itre | | | |
| | | | | | | | | | | | | | | | | | | |
| Doctors' name | | | | | | | | | | | | | | | | | | |
| Name of institution | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | |
| DEMOGRAPHIC 1. Visit date | DE | TΑ | ILS | _ | | 1. | _ | _ | _ | | | | | | | | | |
| (dd/mm/yyyy) | | ╙ | | / | |]/ | | | | | | | | | | | | |
| 2. Name of patient | | | | | | | | | | | | | | | | | | |
| | | N | lyKad | l' Myl | Gd: | | | | | Π. | | | | | Old | HC: | | |
| a uno | | О | ther I | D do | cument N | lo: | | | | | | | | | | | | |
| 3. NRIC | | | pecify pe (if | | ument | ⊚ Regi | | on nu | mber | - | other's U | | - 7 | | rk Permi | _ | | |
| | | " | po (ii | 0.10 | | Pass Birth | | ficate | | - | ather's I/ med For | | | | ver's Lice apital RN | | | |
| | | | | | | | | | | | | | | | | | | |
| SECTION 1 : PA | SIE | EV/ | ۱LU | ΑT | ION | | | | | | | | | | | ı | | |
| Deducation | | | | | | | - | | | eristic | | | | | | Percentage i | involvement of | |
| Body region | | | 0 = N | | . 1 = M | ild, 2 = | | ierat iekn | | = Sew | re, 4 = | | <u>r sev</u> cali | | | | dy region | |
| | | _ | ıyuı | ema | | _ | | CKII | 255 | | - | - | Calli | ıy | | None | ⊚ 50 – 69% | |
| Head | 0 | ⊚ 1 | ⊚ 2 | ⊚ 3 | ⊚ 4 | 0 | ⊚ 1 | ⊚ 2 | ⊚ 3 | ⊚ 4 | 0 | ⊚ 1 | ⊚ 2 | ⊚ 3 | ⊚ 4 | ⊕ 1 - 9% ⊕ 10 - 29% | ⊙ 70 – 89% ⊙ 90 – 100% | |
| | _ | _ | _ | _ | | Ů | _ | _ | _ | | Ľ | | _ | _ | | | | |
| Upper limbs | 0 | 0 | 0 | 0 | • | • | 0 | 0 | 0 | • | • | | 0 | 0 | • | None1 - 9% | ⊚ 50 – 69% ⊚ 70 – 89% | |
| оррег шиоз | 0 | 1 | 2 | 3 | 4 | 0 | 1 | 2 | 3 | 4 | 0 | 1 | 2 | 3 | 4 | ⊙ 10 - 29% ⊙ 30 - 49% | ⊚ 90 – 100% | |
| | 0 | 0 | 0 | 0 | 0 | 0 | • | 0 | 0 | 0 | 0 | 0 | 0 | 0 | • | None1 - 9% | | |
| Trunk | ŏ | ĭ | 2 | 3 | 4 | ő | ĭ | 2 | 3 | 4 | ŏ | | 2 | 3 | 4 | ⊕ 10 - 29% ⊕ 30 - 49% | ⊚ 90 – 100% | |
| | | | | | | | | | | | | | | | | None | ⊚ 50 – 69% | |
| Lower limbs | 0 | 1 | 2 | 3 | ⊚ 4 | 0 | ⊚ 1 | 2 | 3 | ⊙ 4 | 0 | | 2 | 3 | ⊚ 4 | ⊕ 1 - 9% ⊕ 10 - 29% | ⊙ 70 – 89% ⊙ 90 – 100% | |
| | | | | | | | | | | | | | | | | ⊚ 30 – 49% | | |
| SECTION 2 : AD | VE | RSI | E E | VEI | NT(S) | DURII | NG | BIC | LO | GIC | TREA | TME | NT | - | | | | |
| | Adv | | | | | | | | - | | et (dd | | | | | | | |
| ☐ Tuberculosis read | ctival | tion | | | | | | | | | |]/[| | | | | | |
| ☐ Septicaemia | | | | | | | | | | | 1 | / | | | | | | |
| ☐ Lymphoma | | | | | | | | | | | |]/[| | | | | | |
| ☐ Cancer other than | n lyn | npho | ma | | | | | | | | |]/[| | | | | | |
| ☐ Worsening of pso | oriasi | s | | | | | | | Ĺ | | | | | | | | | |
| <u> </u> | | | | | | | | | L | | | | | | | | | |
| <u> </u> | · | | | | | | | | | | | | | | | | | |

| SECTION 3 : BIOLOGIC DOSES RECEIVED | | | | | | | | | | |
|---|---|--|--|--|--|--|--|--|--|--|
| Total number of biologic doses received since initial dose | | | | | | | | | | |
| | | | | | | | | | | |
| SECTION 4 : CHANGE/CESSATION | ON OF BIOLOGIC TREATMENT | | | | | | | | | |
| Date of change/cessation (dd/mm/yyyy) | | | | | | | | | | |
| 2. Change/cessation of biologic treatment | Change to another biologic | | | | | | | | | |
| | Change to another systemic agent. Specify | | | | | | | | | |
| | Withhold biologic treatment | | | | | | | | | |
| | | | | | | | | | | |
| If changed to another biologic agent, please specify | Infliximab IV | | | | | | | | | |
| please specify | Adalimumab SC | | | | | | | | | |
| | Others, specify | | | | | | | | | |
| If changed to another systemic agent, | □ Phototherapy | | | | | | | | | |
| please specify | □ Oral methotrexate | | | | | | | | | |
| | □ Parenteral methotrexate | | | | | | | | | |
| | □ Acitretin | | | | | | | | | |
| | □ Sulphasalazine | | | | | | | | | |
| | · · | | | | | | | | | |
| | Cyclosporin | | | | | | | | | |
| | Hydroxyurea | | | | | | | | | |
| | Systemic corticosteroids | | | | | | | | | |
| 1 | ☐ Other systemic agent, specify | | | | | | | | | |

APPENDIX B: DATA MANAGEMENT

The National Dermatology Registry (DermReg) maintains a database that includes patient's demographic data, medical history, comorbidities, clinical presentation, treatments received in the past 6 months and quality of life. Data is stored in SQL Server due to the high volume of data accumulated throughout the years.

Data Sources

SDPs of DermReg comprise of dermatology centres or clinics with dermatologists who participate in the registry throughout Malaysia.

Data Collection

The study involves collection of data on the patient's first visit to the participating centre and thereafter every six monthly on follow-up visits.

A carefully designed Case Report Form (CRF) is employed in the data collection. This is a double-sided single-sheet CRF which consists of a clinical data form and a multilingual Dermatology Life Quality Index (DLQI) form in both adult and children versions. The clinical data form is to be completed by the doctor in-charge while the DLQI form is to be completed by the patient (parent or guardian for young patient) with guidance from trained staff if necessary. Adult DLQI form should be used for patients above 16 years old, while Children DLQI for patients aged 5 to 16. It is not required to fill the DLQI form for patients below 5 years of age.

One set of CRF is to be completed for each new patient during consultation at the first visit to the participating centre. A new set of CRF is to be completed for the same patient every 6 monthly to record the progress of the patient. The CRFs are used as part of the clinical records.

The CRF is to be completed in duplicate. The participating centre retains the duplicate copy in the patient's medical record, while the original copy is to be sent within 2 weeks to the RCC where data are analysed, interpreted and presented in regular reports to be disseminated to the users.

Participation of SDP is entirely voluntary.


Registry ICT Infrastructure and Data Centre

The operations of the DermReg are supported by an extensive ICT infrastructure to ensure operational efficiency and effectiveness.

The network infrastructure consists of the network layout, placement of relevant hardware equipment, the general flow of data across the network, as well as the network services required for a functional and secure DermReg network infrastructure. DermReg servers are located in a data centre in Cyberjaya in order to provide DermReg with quality assured data hosting services and state-of-the-art physical and logical security features without having to invest in costly data centre setup internally. The physical security features implemented include fire suppression system, access card and biometrics authentication to gain physical access to the data centre, uninterrupted power supply, and backup devices. Logical security features implemented include firewall, antivirus, automated patching, encryption, traffic monitoring and intrusion detection system.


Data Flow Process

Data are collected by doctors in the dermatology departments or clinics. Completed CRFs are then sent to the RCC.



Data received by the RCC are manually reviewed and checked for completeness and error. Data without apparent problems are entered into the registry database. Edit checks are performed periodically to identify potential data errors, such as missing data, non-allowed values, out of range numeric values, inconsistent data and error with deduplication. Data queries that are resolved are then updated to the database.

To ensure complete enumeration and validity of data, a series of tasks as shown in the figure below have to be in place.



SDP Data Reporting, Data Correction and Submission Tracking

Data submitted by SDP are entered into electronic case report form (eCRF) via DermReg Web Application (eDermReg).

There are a number of data security features that are designed into eDermReg such as web owner authentication, two-level user authentication, access control, data encryption, session management to automatically log off the application, audit trail and data backup and disaster recovery plan.

Prior to registering a patient record, a verification process is done by using the search functionality to search if patient exist in the entire registry. This step is done to avoid duplicate records. For patients that exist in the database, SDP only needs to add a new notification with basic patient particulars pre-filled based on existing patient information in the database.

There are a few built-in functionalities at the data entry page that serve to improve data quality. One such function is auto calculation which reduces errors in human calculation. There is also inconsistency check functionality that disables certain fields if these fields are answered in a certain manner. When value entered is not within the specific range, user is prompted for the correct value.

Real time reports are also provided in the web application. The aggregated data reports are presented in the form of tables and graphs manner. These aggregated data reports are typically presented in two manners, one as the centre's own data report and another as registry's overall data report.

Edit checks run and Data cleaning

Edit check was performed periodically by the registry manager to identify missing compulsory data, out of range values, inconsistency data, invalid values and error with de-duplication. Data cleaning is then performed based on the results of edit checks. Data update and data checking of the dataset is performed when there is a query of certain fields when necessary. It could be due to request by user, correction of data based on checking from data query in eCRF or after receiving results from preliminary data analysis. During data standardization, missing data are handled based on derivation from existing data. For example, deriving age from IC, deriving gender from IC and name and inferring race from name. Checking inconsistency of the data also done, for example IC and name shows female but gender is male. Data de-duplication is also performed to identify duplicate records in the database that might have been missed by the SDP.

Legal Aspects and Confidentiality

Data transfer from source data producers is entirely voluntary. There is no legal provision to compel any individual or institution to report or transfer its data to the RCC. The data transferred to RCC is highly sensitive and has to be kept strictly confidential with access only to authorized individual working in the RCC. Strict data protection procedure will need to be put in place, following standard disease registration practice, and in compliance with applicable regulatory guidelines.

Data release policy

One of the primary objectives of the Registry is to make data available to the physicians, policy makers and researchers. The Registry would appreciate that users acknowledge the Registry for the use of the data. Any request for data that requires a computer run must be made in writing (by email, fax, or registered mail) accompanied with a Data Release Application Form and signed Data Release Agreement Form. These requests need prior approval by the Governance Board before data can be released.

Appendix B: Data Management

APPENDIX C: STATISTICAL METHODS

ANALYSIS SET

This refers to the set of cases included in the analysis. Two analysis sets were defined:

1. Patient notification between 2007 and 2011

There were 6,985 patients in the dataset. The analysis set was use for the analysis in Chapter 1, 2, 3, 4, 5 and 6, which comprises of 316 cases in year 2007, 1,283 cases in year 2008, 1,758 cases in year 2009, 1,807 in year 2010 and 1,821 in year 2011. The cases include first notification and up to five follow-up notifications.

2. Patient outcome between 2007 and 2011

There were 1, 936 cases considered for the outcome analysis in Chapter 8.

DATA MANAGEMENT

Data Cleaning

The data from the MPR database were subjected to extensive checking prior to definitive analysis. Errors found or queries raised were checked against the database and/or CRF and corrections were made immediately.

Missing Data

Details on the missing data were issue to Project Manager to clarify the status of the information. Trackable missing information was then incorporated into the dataset but for untrackable and tolerable missing data were included in the analysis and defined as missing.

STATISTICAL METHOD

Descriptive analysis was done in presenting frequencies and percentages of distribution whereas bar and pie charts were used in presenting the figures. For continuous data, the mean, standard deviation, minimum, maximum, median and interquartile range were reported. For standardization in output table, the values of percentages and summary descriptive were limited to one decimal point only. The summaries of data presentation by chapter were described as below:

Stock and Flow

Chapter 1 explained the registry for the distribution of centres reported and distribution of patients according to number of notifications.

Characteristics of Patients

Chapter 2 explained the socio-demographic profiles such as gender, ethnicity, nationality and marital status. Descriptive summary was done for age at visit.

Medical History

Chapter 3 emphasized on the distribution of aggravating factors of psoriasis patients. Crosstabulations were concentrated on the comparison of family members with psoriasis against age of onset.

Comorbidities

Chapter 4 emphasized on the combination of distribution and descriptive summaries of age of onset, several demographic profiles and comorbidities. Figures were presented graphically using bar and stacked bar charts.

Clinical Presentation

Chapter 5 concentrated on the descriptive summaries of pain score. The distribution of psoriasis patients were further analysed on types of psoriasis, body surface area, severity, nail involvement, joint disease, rheumatoid factor, symptoms of psoriatic arthritis and types of joint disease. Crosstabulations performed with several combinations involving age of onset, types of psoriasis, demographic profiles, severities and disease involvements. The graphical presentation were pie chart, bar and stacked bar chart.

Treatment

Chapter 6 presented the distribution of patients with topical therapy, phototherapy, types of phototherapy and systematic therapy. The graphical presenteation were in pie chart, bar and stacked bar chart.

Quality of Life

Chapter 7 solely concentrated on a specific intention, which was on Dermatology Life Quality Index (DLQI). The distribution and crosstabulation figures were presented graphically using bar, stacked bar and line charts.

Outcomes

Chapter 8 explained on the distribution and descriptive summary of the outcome variables. The improvement of lesion extent, skin score, joint score and DLQI score were graphically presented using bar charts.

STATISTICAL SOFTWARE

SPSS 18.0

Appendix C: Statistical Methods

APPENDIX D. PARTICIPATING CENTRE DIRECTORY

| APPENDIX D: PARTICIPAT | ING CENTRE DIRECTORY |
|-----------------------------------|--------------------------------|
| Hospital Kuala Lumpur | Investigator: |
| Department of Dermatology | Dr Azura Mohd Affandi |
| Hospital Kuala Lumpur | |
| Jalan Pahang | Site- coordinator: - |
| 50586 Kuala Lumpur. | |
| Tel: 03-26151540 | |
| Fax: 03-26985927 | |
| Tax. 03-20983927 | |
| Hospital Sungai Buloh | Investigator: |
| Hospital Sungai Bulon | Dr. Azahzuddin b Hamzah |
| Dormetology Unit | DI. Azanzudun o Hanizan |
| Dermatology Unit | Site- coordinator: |
| Hospital Sungai Buloh, | |
| Jalan Hospital, | Prima Dharshini |
| 47000 Sungai Buloh, | |
| Selangor Darul Ehsan. | |
| Tel: 03-61454333 Ext 1286 | |
| Fax: 03-61454222 | |
| Hospital Tuanku Ja'afar, Seremban | Investigator: |
| | Dr. Najeeb Ahmad Mohd Safdar |
| Dermatology Department, | |
| Hospital Tuanku Ja`afar, | Site- coordinator: |
| Jalan Rasah 70300 Seremban, | Dr.Prakash A/L Balasubramaniam |
| Negeri Sembilan. | |
| Tel: 06-760 4157 | |
| Fax: 06-7625771 | |
| | |
| Hospital Sultanah Fatimah, Muar | Investigator: |
| | Dr. Siti Khadijah Abdul Wahid |
| Dermatology Department, | |
| Hospital Pakar Sultanah Fatimah, | Site- coordinator: |
| Jalan Salleh, 84000 Muar, | Mohd Khairul bin Othman |
| Johor. | |
| Tel: 06-9521901 | |
| Fax: 06-9526003 | |
| Hospital Pulau Pinang | Investigator: |
| aropimi i mini i minig | Dr Chan Lee Chin |
| Dermatology Department, | Di Chun Dec Chin |
| Hospital Pulau Pinang, | Site- coordinator: |
| Jalan Residensi, | Dr Yeoh Chin Aun |
| 10990 Pulau Pinang, | Di 100ii Ciiiii Auii |
| Tel: 04-222 5250 Ext 5246 | |
| Fax: 04-2281737 | |
| 1 ax. 04-2201/3/ | |

| Hospital Sultanah Bahiyah, Alor Setar | Investigator: |
|---|--------------------------------------|
| | Dr Mani Mala a/p T. Manikam |
| Dermatology Department, | |
| Hospital Sultanah Bahiyah, | Site- coordinator: |
| Lebuhraya Darul Aman, | Dr Azlida Che Man |
| 05100 Alor Setar, Kedah | |
| Tel: 04-740 6233 | |
| Fax: 04-7350232 | |
| Hospital Tuanku Fauziah, Kangar | Investigator: |
| | Dr Sharifah Farihah Syed Abas, |
| Dermatology Department, | Dr. Hassanin Hussaini Hilmi bin Mohd |
| Hospital Tuanku Fauziah, | Khalid |
| Jalan Kolam,01000 Kangar, | |
| Perlis. | Site- coordinator: |
| Tel: 04-973 8000 | Wan Suhardi bin Wan Abdul Rahman |
| Fax: 04-9767237 | |
| Hospital Queen Elizabeth, Kota Kinabalu | Investigator: |
| | Dr Zaigham Mahmood |
| Dermatology Department, | Dr Mervin George Matthew |
| Hospital Queen Elizabeth, | |
| Karung Berkunci no 2029, | Site- coordinator: |
| 88586 Kota Kinabalu, | Ampong Anggarak |
| Sabah. | 1 6 66 |
| Tel: 088-517555 | |
| Fax: 088-211999 | |
| Hospital Tengku Ampuan Afzan, Kuantan | Investigator: |
| | Dr Abu Razak Yusof |
| Dermatology Department, | |
| Hospital Tengku Ampuan Afzan, | Site- coordinator: |
| Jalan Tanah Putih, | Mazliza binti Mamat |
| 25100 Kuantan, | |
| Pahang. | |
| Tel: 09-5133333 | |
| Fax: 09-5142712 | |
| Hospital Raja Permaisuri Bainun, Ipoh | Investigator: |
| | Dr. Tang Jyh Jong |
| Dermatology Department, | |
| Jabatan Dermatologi, | Site- coordinator: |
| Hospital Ipoh, Jalan Hospital | Dr. Gurcharan Jit Singh |
| 30990 Ipoh, | |
| Perak. | |
| Tel: 05-208 5072 | |
| Fax: 05-2531541 | |
| 1411 00 2001011 | |

| | T |
|---------------------------------------|---|
| Sarawak General Hospital | Investigator: |
| B 1 | Dr Pubalan Muniandy |
| Dermatology Department, | |
| Hospital Umum Sarawak, | Site- coordinator: |
| Jln Tun Ahmad Zaidi Adruce, | Dr Ling Hee Ninh |
| 93586 Kuching, | |
| Sarawak. | |
| Tel: 082-27 6666 Ext 5117 | |
| Fax: 082-242751 | |
| Hospital Tengku Ampuan Rahimah, Klang | Investigator: |
| | Dr. Ng Ting Guan |
| Dermatology Department, | |
| Hospital Tengku Ampuan Rahimah, | Site-coordinator: |
| 41200 Klang, | Dr Norasma bt Roslan, Dr Balachandran a/l |
| Selangor. | Manoharan |
| Tel: 03-3375 7000 Ext 6266 | |
| Fax: 03-3374 9557 | |
| Hospital Sultanah Aminah, Johor Bahru | Investigator: |
| | Dr Choon Siew Eng |
| Jabatan Dermatologi, | |
| Hospital Sultanah Aminah Johor Bahru, | Site- coordinator: |
| Jalan Abu Bakar, 80100, | Hasnal Azahare |
| Johor Bahru, Johor | |
| Tel: 07-223 1806 | |
| Fax: 07-2242694 | |
| Gleneagles Intan Medical Centre | Investigator: |
| | Dr. Chang Choong Chor |
| Hope Skin and Laser Centre, | |
| Gleneagles Intan Medical Centre, | Site- coordinator: - |
| 282 & 286 Jalan Ampang, | |
| 50450 Kuala Lumpur. | |
| Tel: 03-4251 6233 | |
| Fax: 03-4257 9233 | |
| Hospital Melaka | Investigator: |
| | Dr Che Salmi Yusoff |
| Dermatology Department, | |
| Hospital Melaka, | Site- coordinator: |
| Jalan Mufti Hj. Khalil, | Dr Koot Chiew Teen, |
| 75400 Melaka. | Dr. Nor Afalailah Mohd Aris |
| Tel: 06-2892690 | |
| Fax: 06-2841590 | |
| Prince Court Medical Centre | Investigator: |
| | Dr. Gangaram Hemandas Belani |
| Prince Court Medical Centre, | |
| 39, Jalan Kia Peng, | Site- coordinator: - |
| 50450 Kuala Lumpur. | |
| Tel: 03- 26100000 Ext 2955 | |
| Fax: 03-2160 0010 | |
| - | |

| | _ |
|---|--------------------------|
| Universiti Malaya Medical Centre | Investigator: |
| | Dr. Wong Su Ming |
| Department of Medicine, | |
| University of Malaya Medical Centre, | Site- coordinator: |
| Faculty of Medicine, | Dr. Kwan Zhen Li |
| University of Malaya, | |
| 59100 Kuala Lumpur. | |
| Tel: 03-79492429 | |
| Fax: 03-7956 2253 | |
| | T |
| Universiti Kebangsaan Malaysia Medical | Investigator: |
| Centre | Dr. Mazlin Mohd Baseri |
| | a |
| Dermatology Department, | Site- coordinator: - |
| Universiti Kebangsaan Malaysia Medical | |
| Centre, | |
| Jalan Yaacob Latif, | |
| Bandar Tun Razak, | |
| 56000 Kuala Lumpur | |
| Tel: 03-9145 6075 | |
| 03-9145 6640 | |
| Hospital Raja Perempuan Zainab II, Kota | Investigator: |
| Bharu | Dr. Zulrusydi bin Ismail |
| | |
| Dermatology Department, | Site- coordinator: |
| Hospital Raja Perempuan Zainab II, | Muhd. Al-Amin Mat Zin |
| 15586 Kota Bharu, | |
| Kelantan Darul Naim | |
| Tel: 09-7452000 Ext 2384 | |
| Fax: 09-7486951 | |
| 1 47. 07 / 100/31 | |